

3012

HINDUSTAN ANTIBIOTICS

Bulletin


AUGUST 1959

VOL. 2

1



Organic Chemicals



Alcohols	Ketones
Amines	Nitrogen Compounds
Anhydrides	Phenols
Aromatic Hydrocarbons	Plasticisers
Chlorine Compounds	Polyalkylene Glycols
Esters	Rubber Auxiliaries
Glycols	Silicones
Glycol-Ethers	Synthetic Lubricants

Wetting and Emulsifying Agents

INDUSTRIAL PRODUCTS DIVISION
NATIONAL CARBON COMPANY (INDIA) LTD.
BOMBAY • CALCUTTA • DELHI • MADRAS

HINDUSTAN ANTIBIOTICS

Bulletin

Vol. 2

August 1959

No. 1

EDITORIAL COUNCIL

Dr. M. J. Thirumalachar

Chairman & Editor

Shri B. V. Raman

Shri C. N. Chari

Dr. S. R. Sarvotham

Dr. D. Ghosh

Shri A. Neelamegham,

Secretary & Asst. Editor

CONTENTS

Editorial:	
2nd Year	1
Antibiotic Combinations	2
Problems in the Extraction of Penicillin	4
S. R. Sen	
Research Notes:	
S-39, an Antifungal Antibiotic	7
K. S. Gopalkrishnan, K. Chaturbhuj, S. F. Boyce & S. S. Vaidya	
Synthesis of Procaine Hydrochloride	7
R. K. Hulyalkar & P. V. Deshmukh	
N, N'-Dibenzylethylenediamine	9
G. N. Vyas	
Physico-chemical Data on Antibiotics I.	13
A. Neelamegham	
Advertisers' Index	A 4

EDITORIAL & ADVERTISEMENT OFFICES

Hindustan Antibiotics Ltd., Pimpri, Poona Dist.


This *Bulletin* is published on the 15th of February, May, August & November, and contains articles and reviews on all aspects of antibiotics production and use.

The views expressed in this journal are those of the authors and do not necessarily represent those of the Company or of the editors.

Annual subscription : Rs. 3.00 (inland), 8 sh. or \$1.50 (foreign). Single copy: Re. 1.00 (inland), 2sh. 6d. or \$0.50 (foreign).

ADVERTISERS INDEX

Alembic Chemical Works Co. Ltd.	A 13
Bengal Chemicals	A 10
Biddle Sawyer and Co. (India) (P) Ltd.	A 10
Blue Star Engineering Co.	A 11
British Drug Houses (India) (P) Ltd.	A 9
Chika Private Ltd.	A 12
Dumex (P) Ltd.	A 8
Eagle Pharmaceutical Works	A 18
East India Pharmaceutical Works Ltd.	A 17
Franco-Indian United Laboratories	A 5
Geoffrey Manners and Co. (P) Ltd.	A 6
Glaxo Laboratories (India) (P) Ltd.	Inside back cover	A 19
Hoechst Pharmaceuticals (P) Ltd.	A 20
Imperial Chemical Industries	A 18
Indian Process Chemical Laboratories	A 21
Jai Singh's (Dr.) and Son	A 15
Kemp and Co. Ltd.	A 21
Minimax Ltd.	A 16
Mitsui Bussan Kaisha Ltd.	Inside front cover
National Carbon Co. (India) Ltd.	A 4
Oriental Pharmaceutical Industries Ltd.	Outside back cover	A 14
Parry and Co. Ltd.	A 22
Philips India Ltd.	A 24
Protos Engineering Co.	P 3
Sarabhai Chemicals	A 23
Shashi Brothers Private Ltd.	A 7
Swastik Oil Mills	A 9
Swastik Rubber Products Ltd.	
Wyeth (John) and Bro. Ltd.	



With best compliments

from

OPIL


Manufacturers of :

PHARMACEUTICALS & FINE CHEMICALS.

ORIENTAL PHARMACEUTICAL INDUSTRIES LTD.,

64-66, TULSI PIPE ROAD, MAHIM,

BOMBAY 16.



*Maximum Power
for
highest efficacy*

PENIVORAL TRISULFAS

A synergistic combination of
Penicillin V with Sulfadiazine,
Sulfamerazine and Sulfathiazole
for all infections susceptible to
Penicillin and Sulfonamide
therapy.

Vial of 12 tablets

Formula per tablet:

Penicillin V	60 mg. (100000 I.U.)
Sulfadiazine	150 mg.
Sulfamerazine	150 mg.
Sulfathiazole	150 mg.

Also

PENIVORAL FORTE

Vial of 12 tablets
each containing
120 mgs. Penicillin V

PENIVORAL

Vial of 12 tablets
each containing
60 mgs. Penicillin V

For Particulars & Literature

FRANCO INDIAN
UNITED LABORATORIES

Bopar Ghar, Hornby Vellard, Bombay 19.

Serves India

This is our Pledge

To the Country
to maintain the wide distribution which guarantees fair prices in every part of India.

To Consumers
to maintain the high quality of our products and service to your dealers.

To Dealers
to maintain stable prices as far as humanly possible.

We are proud of the reputation we have earned for the first class quality of our products, the fairness of our dealings and our wide distribution and, in this manner, of our service, to India.

GEOFFREY MANNERS & CO. PRIVATE LTD.

Manufacturers of

Anacin • Kolynos • Bismag • Anne French • Aludrox • Endrine • Forhan's • Manners Cough Syrup • Mint Pain Balm • Manners Gripe Mixture • Vaseline Hair Tonic • Wyeth Ethical Products • Manners Ethical Products.

*A decade of experience stands for
faith and confidence with*
"SWASTIK"

A FEW IMPORTANT POINTS WHY

"Swastik"

PRODUCTS HAVE EARNED SO WIDE A POPULARITY

SUPERIOR QUALITY



ATTRACTIVE TERMS



UP-TO-DATE TECHNIQUES OF PRODUCTION



SKILLED SUPERVISION



ECONOMICAL & DURABLE PRODUCTS



THE MARK "SWASTIK" STANDS FOR QUALITY

MAIN LINES OF PRODUCTION :

Electric Cables — Gumboots — Industrial & Surgical Gloves —

Meteorological Balloons — Hose Pipes —

Rubberised Fabrics — Mackintosh & Pure Rubber Sheetings —

Industrial & Auto Parts.


Trade Enquiries Solicited :

SWASTIK RUBBER PRODUCTS, LIMITED

KIRKEE-POONA 3

Phone : 3636

Gram: Swastirub



It takes time to make a doctor

THE ONE LOOK filled with deep gratitude tells more than words can say. Alone it is worth a lifetime of exertion and toil... in the college...in the dispensary...in the hospital...in the home ...a life dedicated to the health and happiness of humanity.

We at Dumex are proud to assist the doctor in his mission with medicines made *in* the country, in modern laboratories, to strict quality specifications laid down in consultation with medical scientists from abroad. Every product is scientifically tested at every stage of manufacture so that you can be certain that every Dumex medicine is pure and fully potent. Dumex medicines are marketed at low prices, too, so as to be of greatest good to the greatest number.

DUMEX

medicines for the people

DUMEX PRIVATE LIMITED, Wavell House, Ballard Estate, Bombay I

DX. 7938

2nd Year....

With this issue, *Hindustan Antibiotics Bulletin*, enters its second year of publication. As befitting the house journal of a national undertaking, the inaugural number was issued on a day of the nation's re-dedication to its people's welfare and progress — 15th August, 1958.

The publication of a journal designed as a medium for fostering better understanding between those engaged in the production and supply of vital drugs and those who use them through dissemination of impartial and authoritative information, obviously entails a good deal of responsibility. Hence, it was with a sense of humility that we placed the first issue in the hands of the scientific and non-scientific public. It is gratifying that our humble effort has been well received all over the world, and judged from the letters of appreciation and encouragement we have received, the journal can embark on the second year with a certain amount of satisfaction.

To all the contributors who placed themselves at our disposal, to the printers and advertisers who extended a co-operative and helping hand, we express our thanks. To the learned societies, institutions and journals who have sent in exchange, valuable scientific and technical publications, and reviewed, abstracted and indexed the *Bulletin*, we are most grateful.

As the journal goes into the second volume we hope to include original articles and other new features in addition to the regular reviews and information sections, so that it may reflect the variety of subjects with which an undertaking such as Hindustan Antibiotics is concerned. We feel assured of the continued support and sympathy of all who have encouraged us and we hope to make many more friends and well-wishers.

EDITORS.

Combined Antibiotic Therapy

Combinations of antibiotics for therapy against human diseases have been widely used. Discovery of a large number of new antibiotics with different antimicrobial spectra, has enabled detailed studies on the combined effects of antibiotics *in vitro* and *in vivo*. Combinations are often used (or misused) when bacteriological data on the disease inciting organism are not available and the patient has to be treated immediately. However, combined therapy has its own advantage as in cases of mixed infection by bacteria, and bacteria and fungi, sensitive to different antibiotics, or in those cases where sufficient antibiotic level cannot be attained by the use of one antibiotic alone. Further, in the case of combined antibiotic therapy, the emergence of resistant organisms is delayed.

When two antibiotics are used as combination in controlling disease incited by a homogenous bacterial population, the effects may be one of the following, as pointed out by Jawetz and Gunnison : (a) *Indifference*—where the combined action is no greater than that of the more potent one when used alone ; (b) *addition*—the combined action is equivalent to the sum of activity of each drug when used alone ; (c) *synergism*—combined action is greater than the sum of activity of both ; (d) *antagonism*—the combined action is less than that of the more effective component when used alone. Jawetz and Gunnison further divided the antibiotics that are in use into two main groups, one comprising the more bactericidal types like penicillin, streptomycin, bacitracin, etc, and the second group including the bacteriostatic ones like tetracyclines, chloramphenicol, and erythromycin.

GROUP I : Penicillin
Streptomycin
Bacitracin
Polymyxin B
Neomycin
Oleandomycin

GROUP II : Chlortetracycline } Tetra-
Oxytetracycline } cyclines
Tetracycline }
Chloramphenicol
Erythromycin
Novobiocin

In therapeutic use, combining members of Group I was considered safe. They would be synergistic or indifferent and never antagonistic. Combining members of Group II would be additive or indifferent and never synergistic or antagonistic. Combinations of Groups I and II might produce antagonism and are rarely synergistic.

In 1943, Ungar was the first to show the synergistic action of penicillin and sulfapyridine, and subsequent clinical experiences with this combination substantiated the original observations. In 1947 Klein and Kimmelman showed synergistic action *in vitro* of penicillin and streptomycin and, this antibiotic pair has been found effective against micrococcic bacterial endocarditis resistant to enormous doses of penicillin. Antibiotic literature is full of reports of cases treated with various antibiotic combinations. Results of the use of penicillin-novobiocin, tetracyclines-oleandomycin, penicillin-streptomycin, tetracyclines-nystatin, tetracycline-novobiocin and others have been described, preference being given by each investigator to some particular antibiotic. In the case of oleandomycin and penicillin, they were even chemically combined as a salt termed "Oleandopen" and shown to have an additive activity.

The concepts synergism, antagonism in combined antibiotic therapy as propounded by Jawetz *et al* have become generally accepted by most of the investigators, but some of the aspects of actual working have often been overlooked. For instance, the influence of bacterial resistance has not been taken into account. If resistance to antibiotics is considered to arise from spontaneous mutations, the number of bacteria present at the time of treatment becomes important. Hence, when there is a large bacterial population, resistant organisms may be present and get selected out by the drug, whereas, in small populations these may be absent.

It is evident that the size of the inoculum influences the final results of antibiotic therapy. Using *Klebsiella pneumoniae*, Ju-

lius and Gaikhorst studied the combined effects of streptomycin and chloramphenicol. When the inoculum potential was low, streptomycin alone was superior to the combined effects of streptomycin and chloramphenicol. This combination would be branded as antagonistic since chloramphenicol reduces the efficacy of streptomycin. However, when the initial inoculum was high, use of streptomycin alone resulted in the rapid development of streptomycin-resistant strains and consequent death of the inoculated animals. Chloramphenicol by itself could not suppress development of resistant strains, but in combination, the resistant strains were suppressed and the action was almost synergistic. It is evident that in this case synergism or antagonism is determined not by the combination of the drugs alone, but is influenced by the amount of initial bacterial inoculum.

INSULATED WIRE & CABLE INDUSTRIES

Ghatkopar—Bombay

Manufacturers of

C. T. S. WIRES & V. I. R. CABLES

Proprietors :

Shashi Brothers Private Ltd.

VASWANI MANSIONS,

DINSHA WACHA ROAD,

BOMBAY I.

Problems in the Extraction of Penicillin

S. R. SEN

The separation and purification of antibiotics from the culture fluids involve a series of critically controlled chemical and physico-chemical operations. Production on a commercial scale consists in starting from an impure biological product obtained by fermentation and ultimately arriving at a chemical with the highest degree of purity having physical properties within strictly set standards and obtaining, at the same time, maximum practicable recovery of the product estimated to be present in the fermented broth. To achieve this object extraction, crystallisation and recrystallisation operations have to be carried out. Ninety per cent of the solids content of the filtered fermented broth are impurities which must be eliminated in order to obtain penicillin crystals of 98 to 99 per cent purity. These impurities interfere with the efficient recovery of the product causing losses and necessitating elaborate handling. The process of extraction has to be so devised as to effect the removal of impurities and increase the concentration of the product in solution from which crystallisation of the solid is practicable. It, therefore, serves as a link between the biochemical process of elaboration of the antibiotic and the chemical process of isolation of the crystals.

Number of variations are feasible in the chemical processes for recovering an antibiotic from its culture fluid. Only a few of them, however, attain commercial acceptance in view of their specific advantages, the percentage recovery obtainable by a particular process being the foremost consideration. Considerations of cost of the chemicals required for processing as well as the hazards in their large scale handling

are also of importance. In general, the recovery processes for antibiotics can be grouped under three heads: Adsorption, solvent extraction and precipitation methods. Often a combination of two or more of these methods are applied to achieve the best results. Carbon adsorption and ion-exchange applications come under the first category. Recovery by these methods are comparatively low and are taken recourse to when it is not possible to recover the product by any other method. Solvent extraction methods involve transfer of the product from one phase to the other thereby eliminating impurities and increasing the concentration in the solvent phase. Precipitation methods are means of isolating the product from its solution as a complex salt and thereafter breaking up the complex to obtain the pure product.

Penicillin is extracted from the fermented broth by the solvent extraction process. In broad general outline the process consists in transferring penicillin from the water solution into an organic solvent phase under acidic conditions. The impurities are partially removed. The penicillin in solution is subjected to another transfer from the solvent to the water phase in the form of its salt. This repetitive process of extraction from solvent phase to water and *vice versa* is carried on in order to obtain a concentration and purity, which makes it possible subsequently to crystallise penicillin by simple operations. In practice two or four extraction steps are employed depending on the conditions prevailing in a particular plant. The quality of the broth and the equipment facilities available determine the yields obtainable by these

two methods. Choice falls naturally on the method giving higher yields.

The transfer operations create certain problems when translated into practice. Most important of these is the inactivation of penicillin. The acid pH required for the efficient transfer of penicillin into the solvent leads to rapid destruction of penicillin in aqueous solutions. In fact, at room temperature the half-life of penicillin in aqueous solution at pH 2 is 18 minutes. Although the rate of inactivation is less at higher pH values, the transfer becomes less efficient. Therefore, to maintain optimum conditions it is essential that the extraction is completed very rapidly. Extraction operation is further complicated by the phenomenon of emulsification. The fermented broth, containing mould metabolites when mixed with the solvent under the acidic condition of extraction, forms emulsion. This is undesirable in that it prevents separation of penicillin-rich solvent phase. Moreover, prolonged and intimate contact of penicillin with the acid accelerates decomposition. This problem is solved by the addition of a surface active agent to the extent of about 0.01 per cent to the broth. It is interesting that wetting agents having emulsifying action are useful in the de-emulsification in this system under the conditions prevailing therein. Again, proteinaceous solids separate out during the process of extraction which interfere with the mixing and separation of the broth and solvent. These solids must be removed continuously in order to facilitate transfer.

The factors controlling the theoretical efficiency of transfer of penicillin from broth to the solvent phase are the distribution coefficient of penicillin between solvent and water at pH 2 and the volumetric ratio of solvent to water. It is obvious, that one stage of mixing and separation will not effect complete transfer unless the distribution is extremely favourable. In the system butyl acetate-water-penicillin a single stage extraction with a solvent to water ratio of

1:4 will give a theoretical efficiency of 83 per cent. A second stage of extraction will bring up the recovery figure to 96.9 per cent, a third stage will theoretically yield 99.3 per cent, a fourth stage 99.87 per cent and so on. This shows that in order to obtain maximum transfer at least three successive stages of mixing and separation are necessary. The process is further improved by employing counter-current extraction. In this system the feed broth solution enters at the end opposite to that of the solvent and the flow of the two phases are one against the other. Thus, the fresh solvent meets partially stripped broth and the enriched outgoing solvents meet the incoming rich broth. The smaller concentration gradients in the stages, therefore, bring about equilibrium conditions quicker. Further, a smaller quantity of solvent is used, thereby attaining greater enrichment.

The foregoing discussion indicates that successful industrial application of penicillin extraction calls for multi-stage counter-current operations with shortest possible time of contact for mixing of phases, rapid separation and effective removal of solids. These particulars should, therefore, be incorporated into the machine selected for the operation. An additional factor is corrosion at the low pH of operation. This can be taken care of by selecting the proper material of construction which usually is molybdenum-stabilized stainless steel.

Rapid technological developments in equipment design met the challenge of this process adequately and today several types of extractors are available which accomplish the transfer of penicillin from the broth to the solvent phase with an efficiency amazingly close to the theoretical.

These equipments must necessarily be of the centrifugal type which would effect very rapid separation, and mixing of the phases is also done centrifugally. Column extractors need longer time of contact for approaching equilibrium conditions and are

therefore of little use in the process. The centrifugal separators develop a relative centrifugal force of the order of 5,000 times gravity or more which is adequate to effect satisfactory separation of two phases having a specific gravity difference as low as 0.02.

The Sharples and DeLaval models of centrifugal extractors are one stage extraction machines. Three stage operation is carried out using three machines with intermediate centrifugal pump mixers. In the DeLaval machine closely spaced set of stationary discs are arranged inside the rotating bowl to increase the speed of separation. The bowl contains adequate space to hold the solids separating out during processing. Models incorporating a device for automatic removal of the accumulated sludge have also been developed, in which as soon as a certain amount of solids separate out the feed is cut-off and a jet of water ejects the sludge out of the machine. Normal operation is continued after discharge of the sludge. This is of definite advantage in that it prevents the tapering off of extraction efficiency due to solid accumulation. Very large capacity machines handling upto 25,000 litres per hour are available. The Luwesta extractor is a similar type of centrifugal separator with discs but with the added feature of having three counter-current stages built-in. Mixing is done inside the machine in small discs with volute shaped passage, utilizing the centripetal force created in the liquid by the rotation of the bowl. One pass through the machine, therefore, effects a three-stage counter-current mixing and separation.

The Podbielniak extractor is a centrifugal machine with strikingly different

features. The rotor consists of a spiral with a rectangular cross section with perforations. The light liquid travels from the periphery to the centre through the spiral path whereas the heavy liquid flows in the opposite direction. The perforations cause intermixing and extraction equivalent to eight theoretical stages is achieved. One particular model of Podbielniak extractor handles unfiltered whole broth. This has proved to be more suited to handling chloramphenicol broth than that of penicillin. Machines having very high throughout capacity of the order of 100,000 litres per hour have been developed. The manufacturers plan to raise this to 200,000 litres per hour in the near future.


It is a fact that the problems in recovering a fermentation product like penicillin have diminished over the past few years due to the advances in fermentation technology. High concentration of a relatively purer product at the end of fermentation, has been achieved. This has made possible higher yields and better quality products with lesser number of processing operations. It is not correct to view extraction processes from the purely chemical standpoint. It must be admitted that the end products of fermentation and the relative purity of the culture fluid will determine the success or otherwise of the extraction process. A high assay of the fermented broth may not necessarily bring about larger output of good quality product. In other words, any effort towards developing the fermentation process must take into account the problems and difficulties likely to be encountered in the separation and purification stages. This alone will make it possible to obtain the final product at the lowest possible cost.

*With Compliments
of*



JOHN WYETH & BROTHER LIMITED

Steelcrete House, Dinshaw Watcha Road
Bombay 1



anacobin

**IN AT THE START
AND STILL LEADING**

Since the inception of Vitamin B₁₂ therapy, 'ANACOBIN' has been synonymous with reliable preparations of pure crystalline vitamin B₁₂. The marketing of 'ANACOBIN' preparations has kept pace with the development of new medical uses for this versatile vitamin—so 'ANACOBIN' is available in a pack and strength for every need.

INJECTIONS: 200 micrograms per ml.
50 micrograms per ml. 500 micrograms per ml.
100 micrograms per ml. 1000 micrograms per ml.

TABLETS:
10 micrograms each
50 micrograms each

ELIXIR:
25 micrograms per
teaspoonful

- A** As a Tonic
- N** Nutritional Macrocytic Anaemias
- A** Anaemias of Pregnancy and Sprue
- C** Cases of Asthma
- O** Osteoarthritis
- B** Bowel disorders of Ulcerative Colitis
- I** In undernourished children for growth promotion
- N** Neuritis of Diabetes

—and also massive doses
for trigeminal neuralgia

BRITISH DRUG HOUSES (INDIA) PRIVATE LTD.


P. O. Box 1341, BOMBAY-1
BRANCHES AT: CALCUTTA • DELHI • MADRAS

With Best Compliments from
BIDDLE SAWYER & CO. (INDIA) PRIVATE LIMITED
 25, Dalal Street, Bombay 1.

Suppliers of :
 Fine Chemicals, Pharmaceuticals,
 Crude Drugs, Specialities,
 Antibiotics, etc. etc.

Agents for :
MESSRS. HENRI DESJONQUERES, Paris.
 Suppliers of Penicillin vials to
 The Hindustan Antibiotics Ltd., Pimpri.

In the treatment
 of *tuberculosis*



• Issued in
 Bottles of 100 & 1000 Tablets.

ISO-NIA-PAS

WITH VITAMINS TABLETS

A powerful chemotherapeutic combination
 for synergistic action in pulmonary and
 other forms of tuberculosis. Counteracts
 development of drug-resistant forms
 of tubercle bacillus.

BENGAL CHEMICAL CALCUTTA • BOMBAY • KANPUR

Extra productivity is
one of many dividends with

Honeywell

Electronik



INSTRUMENTATION

Modern methods of measurement and automatic control can help you raise output by improving operating techniques.

The results—economical operation, increased profits, and high product quality.

Whether you're building an entire new plant, bringing your present plant equipment up to date, or just modernizing a single production unit, Honeywell instrumentation can show the way to greater productivity
...more profits for you.

Honeywell, a pioneer in automatic controls, offers the most comprehensive line in the industry, engineered and serviced in India by BLUE STAR.

Ask a Blue Star representative to advise you on the profitability of modern instrumentation for your plant.



Honeywell *First in Control*

Exclusive Distributors

BLUE STAR



**BLUE STAR ENGINEERING
CO. (Bombay) PRIVATE LTD.**

KASTURI BUILDINGS
JAMSHEDJI TATA ROAD, BOMBAY 1
Also at CALCUTTA, DELHI, MADRAS

7355-98

With the compliments of

Chika
PRIVATE
LIMITED

Suppliers for the products manufactured by



FARBENFABRIKEN BAYER A.G.
LEVERKUSEN — W. GERMANY

- ANTIFOAMS
- SILICONES



CHEMISCHE WERKE HUELS A.G.
MARL — KREIS RECKLINGHAUSEN, W. GERMANY

- ORGANIC SOLVENTS
- ORGANIC ACIDS—
- OTHER ORGANIC CHEMICALS FOR THE
PHARMACEUTICAL INDUSTRIES

Chika **PRIVATE LIMITED**

**Industrial Assurance Building,
Opp. Churchgate Station,
BOMBAY.**

Tel. 2 4 2 2 9 1 - 9 3

CALCUTTA :
31, Chittaranjan Ave.
Tel. 234479

MADRAS :
52/54, Rattan Bazar

DELHI :
Vora House,
Behind Kamala Market

Research Notes

PRELIMINARY NOTE ON S-39, AN ANTIFUNGAL ANTIBIOTIC

In a routine soil screening programme a sample from Darjeeling yielded a species of *Streptomyces*, tentatively identified as belonging to the section of Spira of Pridham *et al*¹, showing strong antifungal properties. The species is considered nearer to *S. albus* than to *S. cacaoi* in spite of some morphological similarities to the latter. *S. cacaoi* has been reported to be only a plant pathogen² and Cacaomycetin which it produces is both antibacterial and antifungal,³ while the species in question produces an antibiotic active exclusively against filamentous fungi and yeast.

Production of the antibiotic in the usual soyabean-dextrose-sodium chloride medium reached a maximum in 96 hrs. giving 250 to 400 *Candida* units per ml. The antibiotic was also produced in other media like glycerol-ammonium sulphate, glucose-ammonium sulphate, and Czapeks, but best production was in soyabean medium. The antibiotic in solution was stable for 4 hr. at 50°, for 1 hr. at 100° and lost 60 per cent activity when autoclaved for 15 min. at 15 lbs. It was active against most of the filamentous fungi tested including *Eremothecium ashbyii*, *Penicillium chrysogenum*, *Aspergillus nidulans*, *Trichothecium roseum*, *Cephalosporium salmosynnematum*, *Fusarium* spp, and *Penicillium herquei*. It was also active against yeasts including *Candida albicans* which was used as a test organism for all bioassays.

The fermentation broth was filtered through Hyflo Supercel and the clear filtrate concentrated to a small volume at 50° under reduced pressure and the con-

centrate extracted with light petroleum (B.P. 40-60°). The extracted broth was acidified to pH 3 and immediately extracted with ethyl acetate, the extract washed with water, dried and the solvent removed to give an oil showing activity against *C. albicans*. The active oil was chromatographed on a mixture of alumina and Hyflo Supercel. Elution with benzene-chloroform (4 : 1) gave an inactive substance, m.p. 147-52°. The benzene-chloroform (1 : 1) fraction yielded an oil which crystallized from ether. Recrystallization from ether gave white crystals, m.p. 97-99°, ultraviolet absorption maxima at 260 m μ ., and soluble in water.

Tested in Sabouraud agar medium, the antibiotic, designated S-39, showed activity against *Candida albicans* at 1:10⁵ dilution.

REFERENCES

1. Pridham, T.G., *et al*. A guide for the classification of Streptomyces according to selected groups. *Appl. Microbiol.* **6**, 52 (1958)
2. Bunting, R. H. Actinomycetes in cacao-beans. *Ann. Appl. Biol.* **19**, 515 (1932)
3. Wakaki, S. Antifungal substances produced by Actinomycetes. II. An antibiotic from a strain resembling to *Streptomyces cacaoi*. *J. Antibiot. (Japan)* **5**, 24 (1952)

K.S. GOPALKRISHNAN, K. CHATURBHUI,
S.F. BOYCE, and S.S. VAIDYA

Plant Mycology Division
and
Organic Chemistry Department
Antibiotics Research Centre
Hindustan Antibiotics Ltd.
Pimpri (Near Poona)

SYNTHESIS OF PROCAINE HYDROCHLORIDE

Based on studies of cocaine degradation and the pharmacological activity of alkylaminobenzoates Einhorn¹ synthesized a series of effective local anaesthetic dialkylaminoalkyl esters. Of a series of alkyl-

p-aminobenzoate type compounds procaine showed the most favourable therapeutic ratio and is now a widely used local anaesthetic. Further, the procaine salt of penicillin was the first successful long-acting penicillin in extensive use even to-day, and hence the commercial importance of procaine.

Procaine has been synthesized by several methods which are subjects of a number of patents. With a view to prepare procaine hydrochloride on a commercial scale we have standardized a process for the preparation of β -diethylaminoethyl-*p*-nitrobenzoate hydrochloride and its direct reduction with Raney nickel. Several reducing agents such as $\text{Zn} + \text{HCl}$ ¹, $\text{Fe} + \text{HCl}$ ², $\text{Fe} + \text{AcOH}$ ^{3,4}, $\text{Sn} + \text{HCl}$ ^{5,2}, $\text{SnCl}_2 + \text{HCl}$ ⁶ and electrolytic reduction⁷ have been used in the earlier methods. Saito *et al*⁸ employed activated nickel for the reduction of the nitro ester in acetone at 80-100° at 20 to 30 atmospheric pressure. In the present work the hydrochloride of the nitro ester in methanol or ethanol was reduced with Raney nickel at room temperature and 50 lbs. pressure, yielding procaine hydrochloride directly in an overall yield of 75 per cent starting from *p*-nitrotoluene. The advantages of the method are that the procaine hydrochloride is obtained directly in one stage by reduction of the corresponding nitro ester hydrochloride and secondly, the reduction of the ester is carried out at room temperature and relatively lower pressure.

EXPERIMENTAL

p-Nitrobenzoic acid⁹

p-Nitrotoluene (250 g.) was oxidized with potassium dichromate (750 g.) and concentrated sulphuric acid (900 ml.) in water (1.5 l.) at 130° for 6 hr. After cooling, ice cold water (3 l.) was added and the precipitate filtered off, dissolved in 10% sodium hydroxide (1.5 l.) and the solution filtered. The filtrate was acidified with 20% sulphuric acid (1 l.),

and the precipitate of pure *p*-nitrobenzoic acid filtered, washed with water and dried; m.p. 236°, yield 250 g. (90%).

p-Nitrobenzoyl Chloride

p-Nitrobenzoic acid (250 g.) was refluxed with thionyl chloride (150 ml.) on a water bath for 16 hr. Thionyl chloride was distilled off and *p*-nitrobenzoyl chloride crystallized from benzene; m.p. 72°, yield 267 g. (98%).

Diethylaminoethyl-*p*-nitrobenzoate Hydrochloride

To *p*-nitrobenzoyl chloride (240 g.) in toluene (1.2 l.) diethylamonethanol (146 ml.) was gradually added and the mixture heated on a water bath for 1 hr. The precipitated hydrochloride of the nitro ester was filtered and washed with toluene (300 ml.); m.p. 127-28°, yield 380 g. (97%).

Procaine Hydrochloride

Diethylaminoethyl-*p*-nitrobenzoate hydrochloride (25 g.) in methanol (150 ml.) was reduced with Raney nickel (0.5 g.) at 50 lbs. pressure and room temperature (28-29°) for 16 hr. Excess of methanol and water was distilled off azeotropically with toluene after removing Raney nickel. The crude procaine hydrochloride was recrystallized from ethanol; m.p. 152°, yield 19 g. (85%).

ACKNOWLEDGEMENT

The authors are thankful to Dr. R. Kaushal for his guidance and Dr. M. J. Thirumalachar for his interest in the work.

REFERENCES

1. Einhorn, A., and Uhlfelder, E. Diethylamino- (and piperido)-ethyl-*p*-aminobenzoate. *Ann.* **371**, 131 (1909)
Idem. Alkamino esters of *m*, *p*-diaminobenzoic acid. *ibid* 162.
Einhorn, A., *et al*. Alkamino ester of *p*-aminobenzoic acid. *ibid* 142.
2. Remington, J. P., Practice of pharmacy. 11th ed., by E. W. Martin and E. F. Cook. Easton, Pa., Mack Publishing Co., 1956, p. 787.

3. Aronovich, I. M., *et al.* Procaine. Russian patent 57, 613 (July 31, 1940). (*Chem. Abst.* **38**, 5047^a, 1944).
4. Nippon Industrial Co. 2-Diethylaminoethyl-p-aminobenzoate hydrochloride. Japanese patent 153, 348 (Oct. 28, 1942). (*Chem. Abst.* **43**, 3847b, 1949).
5. Farbwerke vormals Meister, Lucius & Brüning Höchst a/M. Process of producing p-aminobenzoic acid alkamine esters. German patents 179, 627 (Nov. 27, 1904); 18021 (Jan. 10, 1905), 180292 (Aug. 24, 1905). (*Chem. Abst.* **1**, 2056, 2059, 1907)
6. Einhorn, A. (to Farbwerke vormals Meister, Lucius and Brüning) U. S. patent 812,554 (Feb. 13, 1906). (*Chem. Zent.* **1**, 1365, 1907).
7. Van Winkle, W. A. Novocaine. U. S. patent 1, 501, 635 (July 15, 1923). (*Chem. Abst.* **18**, 2945, 1924)
8. Saito, K., *et al.* (to Nippon Chemical Industries Co.) Alkylamino-ethyl-p-aminobenzoate. Japanese patent 2410 ('51) (May 19). (*Chem. Abst.* **47**, 4913f, 1953)
9. Kamm, O., and Matthews, A. O. p-Nitrobenzoic acid. In Gilman, H., *ed.* Organic synthesis. 2d ed., by A. H. Blatt. New York, John Wiley, 1941. Collected vol. **1**, p. 392.

R. K. HULYALKAR and P. V. DESHMUKH

Antibiotics Research Centre,
Hindustan Antibiotics Ltd.,
Pimpri, Near Poona

N, N'-DIBENZYLETHYLENEDIAMINE*

N, N'-dibenzylethylenediamine (II) is a base for the preparation of N, N'-dibenzylethylenediamine dipenicillin G (Benzathine penicillin), a stable repository penicillin salt; it is also useful for the recovery, purification and preparing formulations of antibiotics.

Bleier¹ prepared (II) from dibenzene-sulphonylethylenediamine; Frost *et al.*², and Menziani *et al.*³, prepared it, along with other by-products, by the conden-

sation of benzylamine with ethylenedihalide; Hauslick *et al.*⁴, alkylated dibenzalidene ethylenediamine with benzylchloride to produce (II). A widely used general method is the reduction of a Schiff's base derived from ethylenediamine and benzaldehyde with sodium and alcohol⁵, or sodium amalgam⁶ and catalytic hydrogenation using platinum oxide⁷, supported palladium⁸, or Raney nickel⁹ at high temperatures and pressures, resulting very often in 1:3 dibenzyl-2-phenyltetrahydroimidazole which is further cleaved to give (II).




(I)



(II)



(III)



(IV)



In the present work, with a view to evolve a simple and economic process for the manufacture of (II), the condensation of ethylenediamine with benzylchloride without the use of alkali¹⁰, has been studied under different conditions. The effects of varying molecular concentrations of the reactants, the period of addition, temperature, and the presence of solvents, on the course of the reaction were investigated to establish optimum conditions for satisfactory yields of (II). The reaction also produced N-mono-benzylethylenediamine (I) in good yields (Cf. Aspinall *et al.*¹¹) which could be converted to (II). The principal advantage of the process is that (II) can be easily separated from the two layers after reaction is complete and directly distilled under reduced pressure without further treatment with water, alkali and extraction with an organic solvent. Further, ethylenediamine, which acts as a reactant as well as an acid binding agent, can be easily recovered for further use. The reaction products mono- (I), di- (II), tri- (III)

* Hindustan Antibiotics (P) Ltd. An improved process for the manufacture of N, N'-dibenzylethylenediamine. Indian Patent 62,439 (Sept. 2, 1958)

and tetrabenzylethylenediamine (IV) were characterized by their hydrochlorides and picrates.

Effect of Molecular Concentration of Reactants

When ethylenediamine was condensed with benzyl chloride in the molecular ratio 1:2, the main product was (IV) (Expt. 1, 2). The molecular concentration of benzylchloride was, therefore, kept constant and that of ethylenediamine gradually increased until the yield of (II) was maximum and that of the higher boiling fractions consisting of (III) and (IV) was minimum. Further increase in the concentration of ethylenediamine led to the formation of (I) in excess of (II) (Expt. 10).

With the molecular concentration ratio of ethylenediamine to benzylchloride at 1:2 or 3:4 the product was mainly (IV) (Expt. 4); at 1:1 ratio a small quantity of (II) and predominantly (IV) were formed (Expt. 5, 6). With 3:2 ratio the results were satisfactory but a fair amount of the higher boiling fractions were also produced (Expt. 7); a ratio of 2:1 was most favourable with a minimum of the higher boiling point fractions. With a 5:1 ratio the main product was (I) with a small amount of (II).

Effect of Solvent

Ethanol and benzene were used as solvents for the reaction. Both the solvents were of no particular advantage when the ratio of the reactants, ethylenediamine to benzylchloride, was 1:2 or 3:4 (Expt. 2, 4). Ethanol improved the

yield of (II) when ratio of reactants was 1:1 (Expt. 5, 6). However, when the ratio was 2:1, ethanol lowered the yield of (II) (Expt. 9).

Period of Addition of Benzylchloride

In small scale experiments the period of addition of benzylchloride was short and of no consequence. But on a large scale, the addition was so regulated to avoid undue rise in temperature leading to formation of higher boiling fractions other than (II) and too slow addition resulting in (I).

Anhydrous ethylenediamine gave better yields of (II) (Expt. 15, 16) but on a larger scale it may be economical to use commercial monohydrate ethylenediamine, there being no appreciable difference in the yield taking into account factors such as the preparation of anhydrous ethylenediamine (Table II).

EXPERIMENTAL

General Procedure

1. Benzylchloride was gradually added to ethylenediamine with shaking and cooling. The mixture was heated on a steam bath for the prescribed period, cooled and treated with sodium hydroxide (10%). (If the solid separated it was filtered and crystallized from alcohol). The filtrate was extracted with ether, the ethereal solution dried over sodium hydroxide, the solvent evaporated and the residue distilled under reduced pressure at 2-3 mm. The different fractions were identified by preparation of the hydrochlorides by passing dry hydrochloric acid in ethereal solution and crystallization from alcohol or acetone.

	B. P./2-3 m.m.	M.P.	
		Hydrochloride	Picrate
Monobenzylethylenediamine (I) 110-15°	253° ¹	221° ¹
N, N'-dibenzylethylenediamine (II) 186-200°	298° ¹	209-11° ¹
Tribenzylethylenediamine (III) 210-30° (m.p. 99°)	Indefinite	Indefinite ²
Tetrabenzylethylenediamine (IV) 235-42 (m.p. 94.5°)	198-202° ¹²	199-200° ¹²

2. Benzylchloride was gradually added to ethylenediamine in alcohol or benzene with shaking and cooling. The reaction mixture was refluxed on steam bath for the prescribed period. If the solid separated it was filtered, treated with dilute alkali (10%) and crystallized from alcohol. The product, m.p. 94-95°, was identified as (IV) as it did not give Hinsberg and Liebermann reactions. The mother liquor was evaporated to remove solvent and further worked up as in 1.

3. Ethylenediamine and benzylchloride were distilled before use. To ethylenediamine monohydrate (78 g., 1. mol.) in a flask kept cool under running tap water, benzylchloride (63.3 g., 0.5 mol.) was gradually added within 20 min. so that the mixture warmed up only slightly. The mixture was heated on steam bath for 12 hr. with continuous stirring, and then transferred to a separatory funnel. The two layers were separated, treated with sodium hydroxide pellets and the liquid fractionated. The lower layer on fractionation gave ethylenediamine, b.p. 118-20°, yield 58 g.; the upper layer on fractionation at 2-3 m.m. pressure gave (I), b.p. 110-15°, yield 15 g.; (II), b.p. 186-200°, yield 25 g. (42% calculated on the amount of benzylchloride used); (III), b.p. 210-30°, yield 3 g. About 3 g. of (II) could be recovered by treating the alkaline solid residue with water, extraction with ether and working up as in 1.

4. N-benzylethylenediamine (15 g.) was mixed with benzylchloride (6.3 g.) and heated on steam bath for 12 hr. The mixture was then treated with sodium hydroxide solution (10%), extracted with ether and dried over sodium hydroxide. The residue, on evaporation of ether, gave (II), b.p. 186-200° /2-3 mm., yield 6 g.

ACKNOWLEDGEMENT

The author wishes to thank Dr. Ganapathi and Dr. M.J. Thirumalachar for their

interest in the work and S. G. Dhopate and A.S. Deshpande for assistance.

REFERENCES

1. Bleier, L. Bildung secundärer Basen aus Aethylen-diamin. *Ber.* **32**, 1825 (1899)
2. Frost, A. E., Jr., et al. Alkyl derivatives of ethylenediamine. *J. Amer. Chem. Soc.* **71**, 3842 (1949)
3. Menziani, E., and Bernabei, M. T. Preparation and analysis of N, N'-dibenzylethylenediamine and its salt with benzylpenicillin. *Bol. Chim. Farm* **93**, 359 (1954)
4. Hauslick, R. S., and Bogash, R. (to American Home Products Corp.) Diaralkylenediamine. U.S. Patent 2,770,653 (Nov. 13, 1956). (*Chem. Abst.* **51**, 8788 (1957))
5. van Alphen, J. Alkylated ethylenediamine derivatives. III. a-Amino-β-benzylaminoethane. *Rec. trav chim.* **54**, 595 (1935)
- Lob, G. 1, 2-Bis (beuzylamino) ethane. *ibid.* **55**, 859 (1936)
6. van Alphen, J. N, N'-Dibenzylethylenediamine. Alkylated ethylenediamine derivatives. I. *ibid.* **54**, 93 (1935)
7. Szabo, J. L. N, N'-Dibenzylethylenediamine penicillin Preparation and properties. *Antibiot. and Chemother.* **1**, 499 (1951)
8. Parker, G (to Distillers Co. Ltd.) N, N'-Dibenzylethylenediamine. Brit. Patent 753,500 (July, 25, 1956). (*Chem. Abst.* **51**, 6690c (1957))
9. Nikiforova, O. K., Preparation of N, N'-dibenzylethylenediamine. *Med. Prom.* **10**, No. 1, 13 (1956). *Chem. Abst.* **51**, 3483b (1957)
10. Jones, W. G. M. (to Imperial Chemical Industries Ltd.) N, N'-Dibenzylethylenediamine. Brit. Patent 762,991 (Dec. 5, 1956). (*Chem. Abst.* **51**, 17991d (1957))
11. Aspinall, S. R. Ethylenediamine. IV. Monoalkyl derivatives. *J. Amer. Chem. Soc.* **63**, 852 (1941)
12. Kerwin, J. F., et al. Adrenergic blocking agents. IV. β-Halo-ethylammonium compounds. *J. Amer. Chem. Soc.* **73**, 5681, (1951)

G. N. VYAS

Antibiotics Research Centre
Hindustan Antibiotics Ltd.
Pimpri, Near Poona

TABLE I
Condensation of Ethylenediamine with Benzylchloride

Expt. No.	Reactants		Solvents		Period of heating on steam bath	Benzylethylenediamine			
	Ethylenediamine monohydrate	Benzyl-chloride	Alcohol	Benzene		Mono-	Di-	Tri-	Tetra-
1.	3.9 g. (0.05 mol)	12.65 g. (0.1 mol.)	—	—	6 hr.	—	—	—	7.0 g.
2.	5.85 g. (0.075 mol)	"	—	50 ml.	5 "	—	—	—	7.8 g.
3.	"	"	—	—	3 hr.	—	—	—	11.2 g.
4.	"	"	40 ml.	—	7 hr.	—	0.7 g.	—	10.2 g.
5.	7.8 g. (0.1 mol.)	"	—	—	5 hr.	—	3.0 g.	—	8.0 g.
6.	"	"	50 ml.	—	12 hr.	—	4.1 g.	2.12 g.	2.1 g.
7.	11.7 g. (0.15 mol.)	"	—	—	6 hr.	2.0 g.	4.5 g.	0.5 g.	—
8.	15.6 g. (0.2 mol)	"	—	—	3.5 hr.	—	3.5 g.	—	—
9.	"	"	60 ml.	—	7.5 hr.	4.0 g.	2.0 g.	—	—
10.	"	5.06 g. (0.04 mol)	—	—					

LARGE SCALE EXPERIMENTS

Condensation of Ethylenediamine with Benzylchloride

Expt. No.	Ethylene-diamine	Benzyl-chloride	Benzyl-chloride addition with tap water	External cooling period	Recovery of Ethylene-diamine	N, N'-Benzylethylenediamine				Remarks
						Mono-(I)	Di-(II)	Tri-(III)	Tetra-(IV)	
11	Monohydrate 78.0 g. (1 mol.)	63.25 g. (0.5 mol.)	20 min.	Yes	58.0 g.	15.0 g.	25.0 g.	3.0 g.		No appreciable diff. without cooling and slow addition.
12	"	"	"	No	62.0 g.	12.0 g.	24.0 g.	4.0 g.		
13	156.0 g. (2 mol.)	126.5 g. (1 mol.)	10 min.	No	134.0 g.	18.5 g.	51.7 g.	12.0 g.	21.0 g.	No cooling, period of addition short; higher yield of (III), (IV)
14	390.0 g. (5 mol.)	316.25 g. (2.5 mol.)	25 min.	Yes	266.0 g.	137.0 g.	138.0 g.	12.0 g.	31.0 g.	Cooling, slow addition; good yields of (I), (II)
15	Anhydrous 36.0 g. (0.6 mol.)	63.25 g. (0.5 mol.)	10 min.	Yes	4.0 g.	8.2 g.	24.0 g.	17.3 g.	4.0 g.	Cooling, period of addition short, amt. of ethylene-diamine small; high yields of (III), (IV)
16	60.0 g. (1 mol.)	"	10 min.	Yes	32.0 g.	14.0 g.	29.0 g.	—	—	Cf. expt. 11; slightly better yield of (II)
17	"	"	35 min.	Yes	28.0 g.	22.0 g.	22.0 g.	6.0 g.	3.0 g.	Same as above, but period of addition long, yield of (I) high
18	"	"	2 min.	Yes	32.0 g.	15.0 g.	29.0 g.	4.0 g.	5.0 g.	Cf. expt. 17; period of addition very short.



"Light of my life..."


"...my pride and my joy", she whispers.

She has watched her son grow from
a tender child to a tall young man.

Your sons, and your sons' sons have all the
necessary care and protection made
possible by modern medical developments.
As a wise patient, do not ignore minor
maladies, consult your doctor. Through
him you will gain the benefit of discoveries
made by the world's scientists. Your
doctor has both the technical knowledge
and the practical experience that will
help you; it's a good habit to

CONSULT YOUR DOCTOR

ALEMBIC CHEMICAL
WORKS COMPANY LIMITED
BARODA-3



Self-Medication can be dangerous ...

... especially for the dyspeptic

Only too often does the chronic hyperchlorhydric take his favourite antacid and aggravate his condition. It is easy to overdose with the usual alkalising treatments and cause acid rebound.

A new compound, introduced here*, cannot cause acid rebound. Its action although rapid, is also *p-r-o-l-o-n-g-e-d* and the stomach-acid is controlled for long periods. It is a buffer-type antacid and hence it is impossible to overdose. It is difficult for the doctor to impress caution on the hyperchlorhydric—the patient will always self-medicate, treat the symptoms himself without advice, varying his own dosage on the wrong assumption that more medicine means more relief.

Agré Antacide promises well in this field. The doctor can advise a safe and effective compound, and the patient has wide dosage-latitude, both qualities leading to possible cure for the chronic sufferer and a more detailed examination of the etiology of the individual's condition.

AGRÉ ANTACIDE

**Aluminium Glycinate*



Sample and literature available from: 'PHILIPS HOUSE', CALCUTTA 20.

PSPH-107

Introducing

Backed with nearly a 100 years experience in the field of pharmaceuticals Kemps offer to the Medical Profession five new products.

Kemvital (ORAL)

A complete balanced Homogenous aqueous multi-vitamin supplementation for all ages.

Kemivit (CAPSULES)

A complete nutritional supplementation providing 11 essential vitamins and 10 important minerals to meet constant need of body tissues.

Kemplivite (CAPSULES)

A miniature Blood bank with all blood building factors a patient needs in anti-anaemic treatment.

Kemp & Co. Ltd.

"Elphin House",

88C, Old Prabhadevi Road,
Bombay-28.

Livokem with B₁₂ (INJECTABLE)

Crude proteolysed Liver Extract containing known and unidentified haemopoietic and nutritional factors, fortified with additional Vitamin B₁₂ for pernicious and other macrocytic anaemias.

Kemplex with B₁₂ (INJECTABLE)

A synergistic formulation of Vitamin B Complex, with Vitamin B₁₂, for intensive treatment in macrocytic anaemias.

KL-R/14.

With Best Compliments from :



MITSUI BUSSAN KAISHA LTD.

(Mitsui & Co. Ltd.)

No. 2, 1-Chome, Shibatamuracho, Minato-ku,
TOKYO (Japan)

WORLD-WIDE IMPORT-EXPORT

Organisation in Japan
With 55 Overseas Offices

New Delhi Branch :
30/48 Panch Sheel Marg
Diplomatic Enclave,
P.O. Box No. 419,
NEW DELHI.

Calcutta Branch :
Suboldutt Bldg.
5th Floor,
13, Brabourne Road,
CALCUTTA I.

Bombay Branch :
Navsari Chambers,
4th Floor,
Outram Road,
Fort,
BOMBAY I.

Telephone Nos. 261667/262148

Cable Address : "MITSUI" BOMBAY

New

CORTOLA-‘m’

Sodium Sulphacetamide—10%

Hydrocortisone—1%

in 3 c.c. dropper bottles

A WORTHY SUCCESSOR TO

CORTOLA

Sodium Sulphacetamide—30%

Hydrocortisone—1%

in 3 c.c. dropper bottles

EAST INDIA



PHARMACEUTICAL WORKS LTD.

CALCUTTA-26

FOR PROMPT CONTROL OF

ALLERGY, INFLAMMATION, TISSUE DAMAGE,
AND ERADICATION OF CAUSATIVE
BACTERIA IN DISEASES OF THE EXTERNAL EYE AND
THE ANTERIOR CHAMBER.

FOR

MERCUROCHROME BPC
POTASSIUM ACETATE BP
SILVER VITELLIN (ARGYROL)
SILVER PROTEINATE (PROTARGOL)
FERRIC AMMON CITRATE BP.
OTHER CITRATES OF
POTASSIUM, SODIUM, CAFFEIN
& SUCH OTHER PHARMACEUTICAL
FINE CHEMICALS OF HIGHEST PURITY

PLEASE CONTACT :

EAGLE PHARMACEUTICAL WORKS

114 Belgrami Road, Match Factory Lane,
KURLA (West), BOMBAY 37

FOR YOUR REQUIREMENTS OF

POTASSIUM ACETATE
POTASSIUM CITRATE
POTASSIUM NITRATE
SODIUM ACETATE
SODIUM CITRATE
BISMUTH CARBONATE
BISMUTH SALICYLATE
BISMUTH SUBNITRATE Etc.

TO APPROVED PHARMACOPOEIAL STANDARDS

PLEASE CONTACT :

INDIAN PROCESS CHEMICAL LABORATORY

YESWANTAPUR P. O., BANGALORE.

Grams: IPCLAB

Phone: 5 5 2 6

Physico-chemical Data on Antibiotics

I. ANTIBIOTICS PRODUCED BY FUNGI, BACTERIA AND LICHENS

1. Melting Points and Ultraviolet Absorption Maxima

Antibiotic activity has been reported for a large number (over 5,000) of microbial culture filtrates and crude substances, but certain amount of chemical, physical and biological data are available only for about 735 antibiotics. As an aid in

Source Organisms	Number of Antibiotics
Bacteria	110
Actinomycetes	305
Fungi and moulds	290
Lichen	30
	<hr/> 735

the identification and study of antibiotics, tables classifying such compounds by physical, chemical and biological data have been compiled. Summaries of the tabulated data will be published in a series of papers. The present compilation is concerned with the physical and chemical properties (melting point, ultraviolet absorption maxima, and empirical formula) of antibiotics produced by bacteria, fungi and moulds, and lichen. Of the 430 antibiotics produced by these groups of organisms, the above data have been reported for about 235 compounds.

In the appended list, the antibiotics are arranged by their empirical formulae, the order being C, H, N, O, Cl, S, other components. The empirical formula being known only for compounds 1-182 the arrangement from serial number 183 onwards is alphabetically by the producing organism. For each antibiotic listed the following data are given: Empirical formula, producing organism, crystal form and colour, nature, melting point, ultraviolet absorption maxima, optical rotation, antibiotic activity and whether the structural formula is known or not.

In Chart 1 for melting points, the serial numbers (S. No.) from the list just mentioned are posted in the squares headed by the relevant melting point range so that S. Nos. of antibiotics having the same

melting point range fall in the same square. For example, antibiotics melting in the range 110-115° are those with S. Nos. 6 and 185, *i.e.* Patulin and Gigantin.

In Chart 2 for ultraviolet maxima, each square represents one wavelength in $m\mu$ on the horizontal side in units (0 to 9) and the numbers (210, 220 etc.) in first vertical column denote the beginning of the range. S. Nos. of antibiotics are posted in the appropriate squares such that those compounds having the same u.v. maxima fall in the same square. For instance, antibiotics with u.v. maxima 235 $m\mu$ are those at S. Nos. 52 and 78 *i.e.* Hydroas-pergillilic acid and Illudin S.

The charts and tests are also helpful in the study of antibiotics as groups of compounds having certain defined chemical-physical properties.

Literature references have been omitted to save space. They are, however, available in comprehensive compilations as:

(1) Spector, W. S., *ed.* Handbook of toxicology. Philadelphia, Saunders, 1957. V. 2 : Antibiotics. (2) Baron, A. L. Handbook of antibiotics. New York, Reinhold Publishing Corp., 1950. (3) Florey, H. W., *et al.* Antibiotics. London, Oxford University Press, 1949. 2V. (4) Korzybski, T., and Kurylowicz, W. Antybiotyki. Warszawa, Panstowowy Zaklad Wydawnictw Lekarskich, 1955. (5) Asahina, Y., and Shibata, S. Chemistry of lichen substances. Ueno, Tokyo, Japan Society for the Promotion of Science, 1954.

ABBREVIATIONS AND SYMBOLS

An asterick Following a S. No. indicates that the structural formula of the compound is known.

A. = *Aspergillus* P. = *Penicillium*

ACKNOWLEDGEMENT

This compilation, originally initiated by Dr. K. Ganapathi, has received the generous help and guidance of the research staff and Dr. M. J. Thirumalachar.

Library, Hindustan Antibiotics Ltd.,
Pimpri, Nr. Poona.

A. NEELAMEGHAN.

CHART 1. MELTING POINT (°C) RANGES

M.P.	25 — 30	30 — 35	35 — 40	40 — 45	45 — 50	50 — 55	55 — 60	60 — 65	65 — 70	70 — 75
S.Nos.		26	91		36	70	29	21 32	33	130 218
M.P.	75 — 80	80 — 85	85 — 90	90 — 95	95 — 100	100 — 105	105 — 110	110 — 115	115 — 120	120 — 125
S.Nos.	144 210	131	21 44 149	11* 38 57	51	8	34 124 125 136	6 185	5 17 121 214	7 123 64 146 78 148 85 209
M.P.	125 — 130	130 — 135	135 — 140	140 — 145	145 — 150	150 — 155	155 — 160	160 — 165	165 — 170	170 — 175
S.Nos.	7 85 120 159	11 54 65 77	22 101 111 134 155	89 99 117	10* 52 88 100	87* 110 129	4 157	47 73 186 220	18* 23 42 53	95 15 75 30 139 37 141
M.P.	175 — 180	180 — 185	185 — 190	190 — 195	195 — 200	200 — 205	205 — 210	210 — 215	215 — 220	220 — 225
S.Nos.	79 98 173	31 113 147 224	115 122 128	96 107 152 158	9* 12* 56	19 60 63* 94	102 107 126	71 76 92 109	72 132 108 115 116	83 206 84* 97 106
M.P.	225 — 230	230 — 235	235 — 240	240 — 245	245 — 250	250 — 255	255 — 260	260 — 265	265 — 270	270 — 275
S.Nos.	28 93 188 35 151 191 55 170* 208 80 172 59 182 92	27 93 184 74 170* 228 80 172 86* 174	48 181	104 176 225	175	20* 163 222*	138 175 192	103*	69 105 175 229	187
M.P.	275 — 280	280 — 285	285 — 290	290 — 295	295 — 300	300 — 305	305 — 310	310 — 315	315 — 320	320 — 325
S.Nos.		193	164	62 162	24	13 171*			14 81	68
M.P.	325 — 330	330 — 335	335 — 340	340 — 345	345 — 350	350 — 355	355 — 360	360 — 365	365 — 370	Above 375
S.Nos.						168	61	196 217		1

* (9) Explodes

(10) Darkens, sublimes

(11) Dec., explodes

(12) Explodes

(18) Chars above 160

(20) Appears to melt at 252

(63) (84) (86) (87) Na salt

(102) Chars on heating

(103) Carbonises

(170) Sulphate 226-28; HCl 332-36 dec.

(171) Above 300

(222) Above 250 dec.

CHART 2. ULTRAVIOLET ABSORPTION MAXIMA

λ_{Max}	0	1	2	3	4	5	6	7	8	9
210	39 41	166 207 208 253	29	7			7 129	70 214		
220	121	26			53	12 69	69 129		70 107 ²	
230	9 77 121 165	165	88 89 111		51 165	52 78	39 53 88 111	41 70	161 215	158 161
240	9 230	18 70 75 116	72 88 100 117	104	111	75	29 53 57 75	26	39 75	41
250	30 40 58 69 98	104 194 215	84 173	178 180 199 200 211		37 183		129	84 85 89 110 113 173 183 ³	60 214 226
260	3 12 35 75 88a	27	13 39 53	31 41 161 227	27 84 173	213 215	110 115	42 110 113 161	73 85 198 201	
270	9 14 56 85	182	216	135 190 212	85 214 226 227	12 130 24 223 ⁴ 32 232 42 91	6 39 85 213	41 197	40 86 202	57
280	69 211 215	129	182		92 93 107	9	70 97 98	83	200	36 129
290	69 142 176 177 214	12 182 226		26	213		129 197	211	30 ¹	

 (1) 297-300 m μ (2) 226-230 m μ (3) 255-260 m μ (4) 270-280 m μ

λ_{Max}	0	1	2	3	4	5	6	7	8	9
300	88 111	5 26		9	117 158	18 42 69 71 72	116	110		214
310	3 12 37 226 227	70		213	4 29 53 88 111	70	75 ¹ 88 111 211	24		
320	69 77 104 110		158 165	9 70 113 115		97 98 121 131	110	57	51 52 53 78 88	111
330	58 69 100	70	13		121		129 158	129	110	211
340	64 69 89			42		20 151				
350	14	158								
360	13 53					196				
370		129							196 223 ²	
380						40				
390	70	70							70	
400	196					3		32 70 130		

(1) 314-318 m μ (2) 375-380 m μ

λ_{max}	0	1	2	3	4	5	6	7	8	9
410										
420	60									
430	70									
440										
450						70 223 ¹				
460										129
470		129								
480										
490										
500						71 72				
510										
520								70		
530						129 ²				129
540		70 129								
550										

(1) 450-460 m μ

(2) 535-540 m μ

S. No.	Empirical formula. Antibiotic.	Produced by	Chemical nature. Melting point (°C)	U. V. absorption maxima (λ max) m μ Optical rotation (α) _D ⁵	Active mainly against (organisms) ⁶
1	2	3	4	5	6
1	C ₅ H ₇ O ₂ Albidin	<i>P. albidum</i>	Glistening red needles. Red pigment. Darkens at 120, no melting at 380.		Fungi. Weak against bac- teria
2	C ₅ H ₆ O ₃ Mycocine C ₁	<i>Penicillium</i> spp. Aceto- bacter spp. <i>Aspergillus</i> spp.			Gram +, gram —, acid fast
3*	C ₆ H ₆ N ₂ O ₄ Toxoflavin	<i>Bacillus cocovenans</i>	Yellow cryst. 172	260, 310, 405 (pH 6.5)	Gram +, gram —
4*	C ₆ H ₆ O ₄ Kojic acid	<i>Acetobacter</i> sp. <i>Aspergillus</i> sp. <i>P. daleae</i>	Colourless prisms. 151-54.	314 (aq. neut. or alk. soln.) Opt. inactive.	Weak against gram + and gram —
5*	C ₇ H ₆ O ₂ Cl. Drosophilin A	<i>Drosophila subatrata</i>	White cryst. Acidic. p-Met- hoxytetraclorophenol, 118 cor.	301. Opt. inactive.	Gram +, <i>M. smegmatis</i> , YSK polio virus, phage Weak against gram —
6*	C ₇ H ₆ O ₄ Patulin	<i>Aspergillus</i> and <i>Penicillium</i> spp.	Colourless rhomboid plates or prisms. Neutral. 109-12.	276. Opt. inactive.	Gram +, gram —, carcinoma cells in tissue culture
7	C ₇ H ₆ O ₄ Terreic acid	<i>A. terreus</i>	Colourless or pale yellow large shining plates. 120-21; 127-127.5.	213, 216. —16 (c, 1 CHCl ₃ , 22°); —28 (c, 1 50% CH ₃ OH— 50% C ₆ H ₆); +74.3 phos- phate buffer, pH 7).	Gram +, gram —, fungi, <i>T. vaginalis</i>
8*	C ₇ H ₈ O ₃ Gentisyl alcohol	<i>Penicillium</i> spp.	Colourless needles. 100-101.		Weak against gram +
9*	C ₈ H ₃ NO ₂ Diatretyne 2	<i>Ciltocybe diatretya</i>	Polyacetylene. 198 uncor., explodes.	230, 240, 270, 285, 303, 323 (95% EtOH).	Activity due to contamina- tion with Diatretyne 1 (12)
10*	C ₈ H ₇ O ₃ Nudic Acid B	<i>Tricholoma nudum</i>	Small colourless needles. 150 darkens, sublimes.	Opt. active in alcohol.	Gram +, gram —
11*	C ₈ H ₅ NO ₂ Agrocycin	<i>Agrocyste dura</i>	White cryst. Neutral, weakly acidic. Ace- tylenic. 90 dec. explodes in vacuo; 130-40 dec. explodes in air.	Opt. inactive.	Gram +, gram —, fungi.

1	2	3	4	5	6
12*	$C_8H_7NO_3$ Diatretyne 1	<i>Clitocybe diatreta</i>	198 uncor., explodes	225, 260, 275, 291, 310 (95% EtOH)	Gram +
13*	$C_8H_8O_5$ Stipitatic acid	<i>Penicillium stipitatum</i>	Cream coloured cryst. Dibasic acid. 302-304 dec.	262, 332, 360 (water) Opt. inactive.	Fungi.
14*	$C_8H_8O_6$ Puberulic acid	<i>P. puberulum</i> , other <i>Penicillium</i> spp.	Colourless microcrystalline plates. Dibasic acid. 316-20 dec.	270, 350.	Gram +
15*	$C_8H_7O_3$ 6-Methylsalicylic acid	<i>P. claviforme</i>	170-71.		Fungi.
16*	$C_8H_7O_3$ 5-Methoxy-p-toluquinone	<i>Lentinus degener</i> , <i>Coprinus similis</i>	Golden yellow plates, clusters of short prisms. Quinone.		Gram +, gram—, acid fast
17*	$C_8H_8O_4$ Fumigatin	<i>A. fumigatus</i>	Maroon needles. Pigmented quinone. 116.		Gram +, gram—
18	$C_8H_7O_5$ Fomecin A	<i>Fomes juniperinus</i>	Cryst. depend on solvent. Weakly acidic. Chars above 160.	241, 305 (95% EtOH). Opt. inactive.	Gram +, influenza virus A. Slight activity against gram—, fungi.
19*	$C_8H_8O_5$ Spinulosin	<i>P. spinulosum</i> <i>A. fumigatus</i>	Purplish bronze or black plates. Quinone. 201-203·5.	Opt. inactive.	Gram +. Lower activity against gram—, acid fast
20	$C_8H_9N_3O_2S$ Micrococin P	<i>Bacillus pumilus</i>	White cryst. solid. Discolours at 232, sinters at 238, appears to melt at 252.	345 (EtOH). +63·7 (c. 1·19 90% EtOH, 21°)	Gram +, <i>M. tuberculosis</i>
21*	$C_7H_{10}O_4$ Penicillic acid	<i>Penicillium</i> spp. <i>Aspergillus</i> spp.	Colourless rhomboid plates or prisms. Monobasic acid. 64-65 (hydrate); 86-87 (anhydrous).	Opt. inactive.	Gram +, gram—, fungi.
22	$C_8H_{10}O_4$ Antibiotic	<i>A. japonicus</i>	134-41 dec. (quinine salt).		<i>Zygosaccharomyces sulcus</i>
23	$C_8H_{13}NO_5$ Oryzasazine	<i>A. oryzae</i>	White hygroscopic needles. 162-63 dec.	—138; —133 (13. 5°)	Sake putrefying bacteria

1	2	3	4	5	6
24*	$C_{10}H_{14}O_7$ Puberulonic acid	<i>P. puberulum</i> , other <i>Penicillium</i> spp.	Yellow plates, prisms, needles. Dibasic acid. 296-98 dec.	274.5, 317	Gram +, less active against gram—.
25	$C_{10}C_8OS_2$ Mycone C_2	<i>Acetobacter</i> spp. <i>Penicillium</i> spp. <i>Aspergillus</i> spp.			Gram +, gram—, acid fast
26*	$C_{10}H_{12}O_2$ 5-Methoxycoumarone	<i>Stereum subpileatum</i>	Colourless, shining leaflets. 34.	221, 247, 293, 300.5 (EtOH)	<i>Staph. aureus</i> , <i>E. coli</i>
27*	$C_{10}H_8O_3$ Flavipin	<i>A. flavipes</i> <i>A. terreus</i>	1: 2-diformyl-4:5:6-trihydroxy-3-methylbenzene. 233-34 dec.	261, 264 (dioxan).	Fungi
28*	$C_{10}H_{15}N_3O_7$ Lycomasramin	<i>Fusarium lycopersici</i>	White microcryst. powder Acidic peptide. 227-29.	—42 to —48 (aq. soln., pH 7).	<i>Lactobacillus casei</i> , <i>Botrytis allii</i>
29*	$C_{10}H_{10}O_3$ Mellein	<i>A. melleus</i> <i>A. ochraceus</i>	Colourless needles. Lactone of 6-hydroxy-2- α -hydroxypropyl-benzoic acid 56-57; 58.	212, 246, 314 (EtOH) —108.15 ($CHCl_3$, 12°)	<i>M. pyogenes</i> var. <i>aureus</i>
30*	$C_{10}H_{10}O_4$ Quadrilineatin	<i>A. quadrilineatus</i>	Colourless needles. Phthalaldehyde. 172 dec.	250, 297-300	<i>Botrytis allii</i> , <i>Pythium debaryanum</i>
31*	$C_{10}H_{12}N_4O_4$ Nebularine	<i>Ciltocybe nebularis</i>	Long prisms, small rhombohedral cryst. Neutral. Furanoside. 181-82-cor.	263 (water). —48.6 (c, 1 water, 25°); —61 (c, 0.8.0.1N NaOH, 25°). —22 (c, 0.8.0.1N HCl, 25°). —47.5 (c, 2 water, 20°). —47.3 (c, 1 water, 20°)	Acid fast. Suppressive effect on ascites tumour in mice
32*	$C_{10}H_{12}O_4$ Aurantio-gluclocladin	<i>Gliocladium roseum</i>	Large orange leaflets. p-Benzquinone. 63.	275, 407.	<i>B. subtilis</i> , <i>Botrytis allii</i>
33*	$C_{10}H_{12}O_4$ Sparasol	<i>Sparassis ramosa</i>	Colourless needles. 67-68		Weak activity on fungi
34*	$C_{10}H_{13}NO_2$ Fusaric acid	<i>Gibberella fujikuroi</i>	Colourless plates. Acid related to nicotinic acid 108-109.		Weakly bacteriostatic

1	2	3	4	5	6
35*	$C_{10}H_{13}N_6O_8$ Cordycepin	<i>Cordyceps militaris</i>	Colourless needles or plates. 22.5-26.	260 (EtOH). —47 (water, 20°)	<i>B. subtilis</i> , <i>Myco. avium</i>
36*	$C_{10}H_{14}O_4$ Glicorosein	<i>Gliocladium roseum</i>	Colourless needles. 48.	289.	Low activity against bacteria
37*	$C_{11}H_7NO_8Cl_2$ Pyoluteorin	<i>Pseudomonas aeruginosa</i>	174-75 dec.	255, 310. Opt. inactive.	Gram +, gram—
38*	$C_{11}H_9O_3$ 6-Methyl-1, 4-naphthoquinone	<i>Marasmius gramineum</i>	Golden yellow needles. 90-91.		<i>M. pyogenes</i> var. <i>aureus</i>
39*	$C_{11}H_9O_3$ Nemotin	<i>Poria corticola</i> , <i>P. tenuis</i>	Neutral. Acetylenic.	207, 236, 248, 262, 276 (water) + 202 (c, 1.2 EtOH); + 350 (alcohol, 17°).	Gram +, gram—, fungi
40*	$C_{11}H_9O_3$ Phthiocol	<i>M. tuberculosis</i>	Bright yellow prisms. Naphthoquinone pigment.	250, 278, 385 (95% alcohol)	Gram +, gram—
41*	$C_{11}H_{10}O_3$ Nemotinic acid	<i>Poria corticola</i> , <i>P. tenuis</i>	Monobasic acid. Acetylenic	208, 237, 249, 263, 277 (water). + 270 (c, 0.44 EtOH, 23°); + 380 (alcohol, 17°).	Same as Nemotin (39), but less active against fungi.
42*	$C_{11}H_{10}O_5$ Gladiolic acid	<i>P. gladioli</i>	Colourless, long, silky needles. Monobasic carboxylic acid. 160.	275, 343 (0.1N NaOH). 267, 305 (Lactol form). Opt. inactive.	Fungi.
43*	$C_{11}H_{10}O_6$ Cyclopaldic acid	<i>P. cyclopium</i>	Colourless, long, fluffy needles. Dibasic acid 224-25.		<i>Botrytis allii</i>
44	$C_{11}H_{15}NO_3$ Candidulin	<i>A. candidus</i>	Long, white needles. Neutral. Non-aromatic. 88-89 uncor.	No characteristic spectrum. + 15 ± 2 (c, 1 CHCl ₃ , 24°).	Gram +, gram—, fungi.
45	$C_{11}H_{15}O_3$ Tardin	<i>P. tardum</i>	Pale yellow oil.	—11.4 (alcohol, 20°)	Gram +, fungi
46	$C_{11}H_{22}O_3$ Pyolipic acid	<i>Pseudomonas aeruginosa</i>	Colourless, odourless, viscous oil. Acidic lipid containing DL- β -hydroxydecanoic acid and a carbohydrate.	—18.9 (20°)	<i>Myco. tuberculosis</i>

1	2	3	4	5	6
47*	$C_{12}H_8N_2O$ Hemipyocyanine	<i>Pseudomonas aeruginosa</i>	Yellow, needles. Pigment, α -hydroxyphenazine. 158.		Gram +, gram—, fungi
48*	$C_{12}H_8N_2O_4$ Iodinin	<i>Chromobacterium iodinum</i>	Purple cryst. with coppery glint. Pigment. 236 dec.		Gram +
49	$C_{12}H_{17}NO_3$ Candidulin (See 44)	<i>Bacillus subtilis</i>	White amorphous powder. Weakly acidic; probably polypeptide.		Gram +, gram—, fungi
50	$C_{12}H_{20}N_2O_2$ Granegilin	<i>Aspergillus</i> sp.	Resembles <i>Aspergillus</i> acid (51).		Gram—, low activity against gram +
51*	$C_{12}H_{20}N_2O_3$ Aspergillic acid	<i>A. flavus</i>	Yellow needles in radial clusters. Monobasic acid. 97-99.	234, 328 +13.4 (c, 0.85EtOH, 24°); +18.5 (c, 1.05 1N NaOH, 24°)	Gram +, gram—, acid fast
52*	$C_{12}H_{20}H_2O_3$ Hydroaspergillic acid	<i>A. flavus</i>	Almost colourless needles. Monobasic acid. 149-50	235, 328. +36 (c, 1.0 EtOH)	Weakly antibacterial
53	$C_{12}H_{20}N_2O_3$ Neohydroxyaspergillic acid	<i>A. sclerotiorum</i>	Colourless, irregular plates. Cyclic hydroxamic acid. 164-66.	236, 328 (EtOH) 246, 360 (0.1N HCl), 224, 262, 314 (hexane)	Gram +, gram—, acid fast, fungi, phage
54*	$C_{13}H_{10}N_2O$ Pyocyanine	<i>Pseudomonas aeruginosa</i>	Dark blue needles. Basic blue pigment. 130 dec.		Gram +, gram—, fungi
55*	$C_{13}H_{10}N_3O$ Chlororaphin	<i>Pseudomonas chlororaphis</i>	Green cryst. Green pigment. 225-30 dec.		Hemolytic streptococci
56*	$C_{13}H_{14}N_2O_4S_2$ Gliotoxin	<i>A. fumigatus</i> , other fungi	Colourless, long plates or needles. 195.	270. —239 — to —256 (CHCl ₃); —290 +10 (0.078 EtOH, 25°)	Gram +, gram—, fungi, sarcoma implants

1	2	3	4	5	6
57	$C_{13}H_{14}O_3$ Diaporthin	<i>Endothia parasitica</i>	Colourless cryst. 91.5-92.5 cor.	246, 279, 327 +58 (c, 1 $CHCl_3$)	<i>B. subtilis</i> , fungi
58*	$C_{13}H_{14}O_5$ Citrinin	<i>P. citrinum</i> , other <i>Penicillium</i> and <i>Aspergillus</i> spp.	Golden prismatic needles or serrate plates. Monobasic acid. 170, 175 dec.	250, 330 (∞) ₅₄₆₁ —43.1 (c, 0.92 absol. EtOH);—37.4 (c, 1.15 EtOH, 18°)	Gram +
59*	$C_{14}H_{10}O_4$ Phenicin	<i>P. phoenicum</i> , <i>P. rubrum</i>	Yellow-brown, long rectangular tablets. Dibasic acid pigment. 229-31.		Gram +, <i>Myco. phlei</i>
60*	$C_{14}H_{10}O_4Cl_2$ Mollisin	<i>Mollisia caesia</i> , <i>M. fallens</i>	Orange yellow needles. 202-203.	259, 280sh, 420, (5.4 mg. in 0.5 mol. $CHCl_3$) Opt. inactive.	Fungi
61*	$C_{14}H_{10}O_5$ Alternariol	<i>Alternaria tenuis</i>	Colourless needles Substituted dibenzo- α -pyrone. 350 dec.		<i>M. pyogenes</i> var. <i>aureus</i> <i>E. coli</i>
62*	$C_{14}H_{10}O_7 \cdot 2H_2O$ Citromycetin	<i>P. frequentans</i>	Yellow cryst. Dibasic acid. 290-300 dec.	Opt. inactive.	<i>M. pyogenes</i> var. <i>aureus</i>
63*	$C_{14}H_{20}N_2O_4S$ Pentenyl penicillin (Penicillin F)	<i>P. notatum</i> , <i>Penicillium</i> spp., <i>Aspergillus</i> spp.	Colourless prisms. Monobasic carboxylic acid. 204-205 dec. (Na salt)	No characteristic bands. +276 to +316 (c, 0.821 water, 20-25°).	Gram +, spp. of <i>Treponema</i> , <i>Borrelia</i> , <i>Leptospira</i>
64	$C_{14}H_{20}O_3$ Nadic acid A	<i>Tricholoma nudum</i>	Colourless needles or plates. 123.5 uncor.	Active in alcohol.	Gram +, gram—
65	$C_{14}H_{20}O_4$ Frequentin	<i>P. frequentans</i>	Colourless needles. Aldehyde. 134.5 dec.	+82 ($CHCl_3$, 24°)	Fungi, low activity against bacteria
66*	$C_{14}H_{21}N_9O_6S$ Synnematin B; Cephalosporin N	<i>Cephalosporium salmosynematum</i>	Ba salt: White powder. Hydrophilic penicillin, containing D- α -aminoadipic acid.	Ba salt: +187 (c, 0.6 water, 20°).	Gram +, gram—. <i>Myco. tuberculosis</i> at higher conc.
67*	$C_{14}H_{25}O_4$ Paltantin	<i>P. paltans</i>	Colourless needles. Unsaturated dihydroxy aldehyde (Cf. frequentin), 163.	(∞) ₅₄₆₁ : +4.4 (c, 0.8, $CHCl_3$).	

1	2	3	4	5	6
68*	$C_{15}H_{10}O_5$ Strepsilin	<i>Cladonia streptilis</i>	Long prisms or needles. Dibenzofuran derivative. 324.		<i>Staph. aureus</i> , <i>Myco. tuberculosis</i>
69	$C_{15}H_{11}NO_2$ Viridicatin	<i>P. viridicatum</i> . <i>P. cyclopium</i>	Colourless cryst.; lustrous needles (from EtOH); prisms (from other solvents) 269	225, 280, 305, 320, 330 (EtOH) 226, 250, 290, 330, 340 (a.q. NaOH). No rotation	<i>Myco. tuberculosis</i>
70*	$C_{15}H_{14}O$ Lactaroviolin	<i>Lactarius deliciosus</i>	Violet red pigment. Azulenealdehyde. 53	241, 311, 390, 541 (cyclohexane). 242, 315, 398, 527 (alcohol). 286, 331, 407, 455 (formic acid). 217, 228, 237, 323, 391, 430 (50% H_2SO_4)	<i>Myco. tuberculosis</i>
71*	$C_{15}H_{14}O_6$ Javanicin	<i>Fusarium javanicum</i>	Red laths with coppery lustre Weak monobasic acid. 207.5-208 dec.	305, 505 (EtOH) Opt. inactive.	Gram +, acid fast
72*	$C_{15}H_{14}O_7$ Oxyjavanicin	<i>Fusarium javanicum</i>	Red needles with coppery lustre 212; 213-16 dec.	305, 505 (EtOH) Opt. inactive.	Gram +, acid fast
73*	$C_{15}H_{16}N_2O_5S_2$ Gliotoxin monoacetate	<i>P. terlikowski</i>	Pale yellow rhombic cryst. 159-60	268 —197 (c, 0.6 $CHCl_3$, 19°)	Same as for Gliotoxin (56) but only one-tenth activity
74*	$C_{15}H_{16}O_5$ Fuscin	<i>Oidiodendron fuscum</i>	Orange, diamond-shaped plates. Quinonoid pigment. 230	Opt. inactive.	Gram +, low activity against gram—
75	$C_{15}H_{18}O_4$ Marasmic acid	<i>Marasmius conigenus</i>	White, long needles. Monobasic acid. 174-75 cor. (in sealed capillary in vacuo).	240-242, 314-318 (EtOH). 245, 246, 248, 260 (phosphate buffer, pH 1.45-3.52, 5.4, 8.8, 11.0). +176 (c, 1.4 C_3H_6O , 25°)	Gram +, acid fast, less active against gram—. Fungi.
76*	$C_{15}H_{18}O_5$ Dihydrofuscin	<i>Oidiodendron fuscum</i>	Colourless rhombic cryst. Quinol. 206 dec.	Opt. inactive.	Gram +, gram—
77	$C_{15}H_{20}O_3$ Illudin M	<i>Ciltocybe illudens</i>	Neutral 130-31 uncor.	230, 320, (EtOH). —126 (absol. EtOH, 20°).	Acid fast. Weak antifungal activity

1	2	3	4	5	6
78	$C_{15}H_{20}O_4$ Illudin S	Ciltocybe illudens	Neutral 124-25	235, 328 (EtOH) —165 (absol. EtOH, 20°).	Gram +, acid fast
79	$C_{15}H_{20}O_4$ Hirsutic acid C	Stereum hirsutum	Colourless, large prisms 179.5 uncor.	None. +11.9 (free acid in absol. alcohol, 20°)	Inactive precursor of active acids.
80*	$C_{16}H_{14}O_5Cl_4$ Diploicin	Buellia canescens	Colourless coral form cryst. Despidone of orcinol group. 232		C. diphtheriae mitis, acid fast.
81*	$C_{16}H_{10}O_7$ Endocrocin	A. amstelodami Nephronopsis endocrocea	Orange red cryst. Quinone 318		Staph. aureus
82	$C_{16}H_{14}O_6Cl_2$ Estin	P. paxilli var. echinula- tum	223-25		M. pyogenes var. aureus, Myco. tuberculosis
83	$C_{16}H_{17}N_3O_4S_2$ Cheomoin	Chaetomium spp.	White non-cryst. powder. Neutral? 215; 218-20 dec.	287. +360 (c, 1 CHCl ₃ , 22°).	Gram +, acid fast
84*	$C_{16}H_{18}N_2O_4S$ Benzylpenicillin (Penicil- lin G)	P. notatum, other Penicil- lium, Aspergillus spp.	Colourless prisms Monobasic carboxylic acid 215 dec. (Na salt)	252, 257.5, 264 +305 (c, 0.821 water, 25°); +301 (c, 2 water, 25°)	Gram +, spp. of Treponema. Borrelia, Leptospira
85*	$C_{16}H_{18}N_2O_5S$ Phenoxymethyl penicillin (Penicillin V)	As for penicillin G (84)	Colourless prisms Monobasic carboxylic acid 120-28	268, 274 (water) 270, 276 (CHCl ₃)	As for penicillin G (84)
86*	$C_{16}H_{18}N_2O_5S$ p-Hydroxybenzyl peni- cillin (Penicillin X)	As for penicillin G (84)	Colourless prisms Monobasic carboxylic acid 228-35 dec. (Na salt)	278. +267 (c, 0.525 water).	As for penicillin G (84)
87*	$C_{16}H_{18}N_2O_6S$ p-Hydroxyphenoxy- methyl penicillin	P. chrysogenum	Colourless prisms 151-59 dec. (Na salt)	+215 (c, 1.06 Na salt in water)	As for penicillin G. (84)
88	$C_{16}H_{21}NO$ Pyo Ib	Pseudomonas aeruginosa	Colourless cryst. 2-Heptyl-4-quinolinol 146.2-147	236, 316, 328 (95% EtOH); 232, 300 (ethanolic HCl); 242, 314 (ethanolic KOH)	
88a	$C_{16}H_{21}N_3O_8S$ Cephalosporin C	Cephalosporium sp.	Na salts: Monoclinic cryst. Hydrophilic penicillin con- taining D- α -amino-adipic acid	260 +103 (c, 0.9 water, 20°)	Staph. aureus, Salm. typhi (1/10 activity of Cephalo- sporin N (66))

1	2	3	4	5	6
89	$C_6H_{23}NO_3$ Pyo IV	<i>Pseudomonas aeruginosa</i>	Colourless needles 131-32; 139, 5-140	232, 258, 340 (95% EtOH)	
90*	$C_{16}H_{28}N_2O_4S$ n-Heptyl penicillin (Penicillin K)	As for penicillin G (84)	Colourless prisms Monobasic carboxylic acid	No characteristic spectrum. +258 (c, 0.43 water, 25°)	As for penicillin G (84)
91	$C_{16}H_{28}O_2$ Grifolin	<i>Grifola confluens</i>	Colourless fine needles 40	275. Complete transmission above 300	Gram +, less active against gram- and acid fast
92	$C_{17}H_{19}O_7Cl_3$ Geodin-like antibiotic	<i>A. flavipes</i>	Pale yellow elongated micro- prisms Dibasic acid 208 (form I), 229-30 (form II)	284. +175 (CHCl ₃ , 20°)	Gram +, Myco. pseudotuber- culosis
93*	$C_{17}H_{12}O_7Cl_3$ Geodin	<i>A. terreus</i>	Brown or yellow needles 227-30; 235 dec.	284. (∞) ₅₄₆₁ ²⁰ +179 (c, 0.8 CHCl ₃); ₅₇₈₀ ²⁰ +149 (c, 0.8 CHCl ₃).	Gram +, gram-, Myco. smegmatis
94	$C_{17}H_{14}O_4$ Canescin	<i>P. canescens</i>	Colourless needles (from EtOH) Monobasic acid 201-202 dec.		Fungi. Low antibacterial activity.
95*	$C_{17}H_{16}O_7$ Evermic acid	<i>Evernia prunastri</i> , <i>Usnea</i> sp., <i>Ramalina pollinaria</i>	Prisms Depside of orcinol group 169-70		Myco. tuberculosis
96	$C_{17}H_{16}O_7$ Antibiotic	<i>Monosporium bonorden</i>	Colourless cryst. 193-5	+203 (CHCl ₃ , 20°).	Fungi.
97*	$C_{17}H_{17}O_6Cl$ Griseofulvin	<i>P. griseofulvum</i> , other <i>Penicillium</i> spp.	Colourless, massive, rhombic or octahedral cryst. Neutral 218-221	286, 325 +370 (c, 2 CHCl ₃ , 17°); +337 (c, 1 C ₃ H ₆ O, 21°)	Fungi
98*	$C_{17}H_{18}O_6$ Dechlorogriseofulvin	<i>P. nigricans</i>	Needles Neutral 179-81	250, 286, 325 +390 (c, 1 C ₃ H ₆ O, 19°)	<i>Botrytis allii</i>
99*	$C_{17}H_{20}O_6$ Mycophenolic acid	<i>P. brevicompactum</i> , other <i>Penicillium</i> spp.	Colourless radiating needles Weak dibasic acid 141	Opt. inactive	Gram +, fungi

1	2	3	4	5	6
100	$C_{17}H_{23}NO_2$ Pyo II	<i>Pseudomonas aeruginosa</i>	Light yellow platelets 149-149.5	242, 330 (95% EtOH)	Gram+, gram— less sensitive
101*	$C_{17}H_{23}O_4$ Roccellic acid	<i>Rocella</i> spp., <i>Lecanora</i> spp.	Plates or leaflets Dibasic acid 130, 131		Myco. tuberculosis t. bovinus
102*	$C_{18}H_{12}N_2O_2$ Xanthocillin X	<i>P. notatum</i>	Yellow clusters of needles (from alcohol); yellow rhombic prisms (from ethyl acetate) Chars at 200		Gram+, gram—, Myco. tuberculosis
103*	$C_{18}H_{12}O_{10}$ Salazinic acid	<i>Parmelia</i> spp.	Needles Depsidone of β -orcinol group Darkens from 240 and carbonises at 260		Staph. aureus
104	$C_8H_4O_7$ Bostrycoidin	<i>Fusarium bostrycoides</i>	Brown or red laths in clusters. Substituted naphthoquinone pigment 243-44	251, 320	Gram+, Myco. tuberculosis
105*	$C_{18}H_{14}O_8$ Psoromic acid	<i>Psoroma crassum</i> , other lichen spp.	Needles. Depsidone of β -orcinol group 265		Staph. aureus
106*	$C_{18}H_{15}O_9Cl$ Pannarin	<i>Pannaria</i> spp.	Long prisms Depsidone of β -orcinol group 216-17		Staph. aureus
107*	$C_{18}H_{16}O_7$ Usnic acid	Lichen spp., fungal symbiont of lichen	Yellow cryst., form differs with solvent of crystallization (c, 2.0-3.19 $CHCl_3$, 20°) 203-204 (d-form); 203 (l-form); 194 (dl-form).	226-230, 284 D-isomer: +445 to +515 (c, 2.0-3.19 $CHCl_3$, 20°) L-isomer: —445 to —478 (c, 2.0-3.08 $CHCl_3$, 20°)	Gram+, gram—, acid fast, fungi.
108	$C_{18}H_{16}O_8Cl_2$ Nordin	<i>P. paxilla</i> var. <i>echinulatum</i>	134-35. Contaminated with Estin has m.p. 214-16.		
109	$C_{18}H_{18}O_7$ Obtusatic acid	<i>Ramalina</i> spp.	Needles. Depsidone of orcinol group 208-209		Staph. aureus

1	2	3	4	5	6
110	$C_{18}H_{23}NO$ Pyo III	<i>Pseudomonas aeruginosa</i>	Colourless cryst. 2-(Δ' -nonenyl-)-4-quinol- inol 151.5-153.5.	258, 266, 307, 338 (95% EtOH). 258, 267, 320 (ethanolic HCl); 260, 326 (ethanolic KOH)	Gram, + gram—
111	$C_{18}H_{25}NO$ Pyo Ic	<i>Pseudomonas aeruginosa</i>	2-Nonyl-4-quinolinol. 138.8-139.2.	236, 316, 328 (95% EtOH); 232, 300 (ethanolic HCl); 244, 314 (ethanolic KOH)	Weak antibacterial activity.
112	$C_{18}H_{32}O_2$ Linoleic acid	<i>P. crustosum</i>			
113*	$C_{19}H_{14}O_4Cl_3$ Nidulin	<i>A. nidulans</i>	Colourless rhombs (from EtOH); shining slender rods (from petrol ether) 180	267, 323 inf. Opt. inactive ($CHCl_3$)	Myco. tuberculosis
114*	$C_{19}H_{14}O_5$ Vulpinic acid	<i>Evernia</i> spp. <i>Cetraria</i> spp.	Yellow prisms (from ether or EtOH); Yellow plates (from from C_6H_6 or $CHCl_3$); Aromatic acid. 148		Myco. tuberculosis avium
115*	$C_{19}H_{15}O_5Cl_3$ Normidulin	<i>A. nidulans</i>	Heavy square plates (fraction I); rosettes of fine needles (fraction II); hexagonal plates or prisms (from pet- rol ether). 185-87 (fraction I); 214-16 (fraction II).	266, 323 inf.	Gram +, acid fast.
116	$C_{19}H_{16}O_6$ α -Viridin	<i>Trichoderma viride</i>	Colourless fine needles 208-17 dec.; 217-23 dec. (α , β mixture)	241.3, 304.5 (alcohol) —213.4 ($CHCl_3$, 20°)	Fungi. Weak activity against protozoa
117	$C_{19}H_{16}O_6$ β -Viridin	<i>Trichoderma viride</i>	Fine needles. 140-dec. 217-23 dec. (α , β mixture)	241.5, 304 (alcohol) —50.7 ($CHCl_3$, 20°) Viridins: —222 (c, 1 $CHCl_3$, 19°)	Fungi. Weak activity against protozoa
118*	$C_{19}H_{16}O_{11}$ Thamnolic acid	<i>Thamnomia</i> <i>vermicularis</i> , <i>Cladonia</i> spp., other lichen spp.	Faint yellow plates. Depside of β -orecinol group. 223 dec.		Staph. aureus
119*	$C_{19}H_{18}O_8$ Atranorin	<i>Evernia prunastri</i> , other lichen spp.	Colourless prisms. Depside of β -orecinol group. 196		Staph. aureus

1	2	3	4	5	6
120	$C_{19}H_{20}O_8$ Herquein	<i>P. herquei</i>	Yellow-brown cryst. 129 dec.		Gram+, gram—, Myco. phlei
121	$C_{19}H_{24}O_6$ Trichothecin	<i>Trichothecium roseum</i>	Colourless, slender, fibrous needles. Neutral ester. 118 cor.	220, 334 ($CHCl_3$ or hexane); 230, 325 (EtOH) +44 (c, 1.0 $CHCl_3$, 18°)	Fungi
122	$C_{19}H_{26}O_8$ Rosein II	<i>Trichothecium roseum</i>	Colourless fibrous needles. 186	+5.9 (c, 2 $CHCl_3$, 23°); (∞) 540.1 +7.5 (c, 1 EtOH, 20°)	<i>B. subtilis</i> , Myco. phlei
123*	$C_{19}H_{26}O_4$ 1-Licheterinic acid	<i>Cetraria</i> spp.	Needles. Monobasic lactic acid. 124.		<i>Staph. aureus</i> , Myco. tuber- culosis avium
124*	$C_{19}H_{32}O_4$ d-Protolicheterinic acid	<i>Cetraria islandica</i> , <i>Parme- lia simodensis</i> , <i>Cladonia</i> spp.	Colourless, rhombic, leaflets with pearly lustre. Monobasic lactic acid. 107.5.		<i>Staph. aureus</i>
125*	$C_{19}H_{32}O_4$ 1-Protolicheterinic acid	<i>Cetraria crispa</i>	Leaflets with pearly lustre. 106.		<i>Staph. aureus</i>
126*	$C_{20}H_{16}O_6$ Pimaric acid	<i>Cetraria</i> spp., <i>Lepraria flava</i> f. <i>quercina</i>	Orange needles. Related to phenylbenzoqui- none pigment. 200-203.		<i>Staph. aureus</i>
127*	$C_{20}H_{17}O_5Cl_3$ Nidulin (See 113)	<i>Pleurotus griseus</i>	Yellow, orange, amber needles. Neutral. 200-15 dec.	250. —20 (c. 0.59 $CHCl_3$, 23°)	Gram+, acid fast. Gram— less sensitive
128*	$C_{20}H_{22}O_7$ Diffractaic acid	<i>Usnea diffracta</i> ; <i>Alectoria ochroleuca</i>	Needles. Depside of β -orcinol group. 189-90.		<i>Staph. aureus</i>
129*	$C_{20}H_{26}N_3O$ Prodigiosin	<i>Serratia marcescens</i>	Green, metallic leaflets; lus- trous square pyramids; dark green with green reflex pigment. Red 151-52.	535-540 (acid-chloroform); base in EtOH; pH 7.4; 225.5, 288.5, 337, 471, 539; pH 11: 257, 281, 335.5, 468.5; pH 2.9: 216, 296, 371, 541 with inflec. at 275, 382, 510	Fungi and protozoa

1	2	3	4	5	6
130*	$C_{20}H_{28}O_8$ Rubrogliocladin	Gliocladium roseum	Dark red needles. Quinhydrone. 74.	275, 407	Botrytis alli; less active against <i>B. subtilis</i> , <i>E. coli</i>
131	$C_{20}H_{32}O_8$ Oregonensin	Ganoderma oregonense	White needles. Neutral. 82.	325 Opt. inactive	<i>M. pyogenes</i> var <i>aureus</i> , gram — and acid fast less sensitive
132	$C_{21}H_{18}O_6Cl_2$ Compound II	<i>A. nidulans</i>	214-16.		Gram + cocci and Myco- bacteria in synthetic media
133*	$C_{21}H_{28}O_5Cl$ Sclerotiorin	<i>P. sclerotiorum</i> , <i>P. multi-</i> <i>color</i>	Yellow cryst. Pigment. Azaphilone. 205-206.	+500 (c. 1 $CHCl_3$, 21°); (∞) 5790 + 549 (21°)	Slight antibacterial activity
134*	$C_{21}H_{24}O_4$ Divaricatic acid	<i>Evernia</i> spp.	Needles. Depside of orcinol group. 137.		<i>Staph. aureus</i> , Myco. tuber- culosis avium
135	$C_{21}H_{30}O_8$ Alternaric acid	<i>Alternaria solani</i>	Colourless, thin rectangular plates or elongated prisms. Unsaturated dibasic acid. 138.	273. Opt. inactive	Fungi
136*	$C_{21}H_{32}O_6$ Rangiformic acid	<i>Cladonia mitis</i> , <i>C. rangi-</i> <i>formis</i>	Needles. Tribasic acid. 106.		<i>Staph. aureus</i>
137*	$C_{21}H_{32}O_7$ Caperatic acid	<i>Parmelia caperata</i> , <i>Nephromopsis</i> sp.	Leaflets. Tribasic acid. 132-133.5.		<i>Staph. aureus</i>
138*	$C_{22}H_{16}O_{12}$ Fumarprotocetraric acid	<i>Cetraria islandica</i> , <i>Clado-</i> <i>nia</i> spp.	Small needles. Depsidone of β -orcinol group. 250-60.		<i>Staph. aureus</i>
139*	$C_{22}H_{26}O_5$ Didymic acid	<i>Cladonia</i> spp.	Needles. Dibenzofuran derivative. 172-73.		<i>Staph. aureus</i> , Myco. tuber- culosis avium
140*	$C_{22}H_{28}O_8$ Sekikaic acid	<i>Ramalina</i> spp.	Needles. Depside of orcinol group. 147 dec.		<i>Staph. aureus</i> , Myco. tuber- culosis avium

1	2	3	4	5	6
141*	$C_{22}H_{28}N_2O_5$ Myceliamide	<i>P. griseofulvum</i>	Colourless shiny leaflets. Amide of o-mycetyl-N-pyrro- vyl- β -ketotyrosine. 170-72.	$(\infty)_{540}:-217$ (c, 0.8688 CHCl ₃ , 19°); $(\infty)_{570}:-182$ (c, 0.688 CHCl ₃ , 19°)	Gram +
142	$C_{22}H_{34}O_5$ Pleuromutilin (Droso- philin B)	<i>Pleurotus mutilis</i> , <i>Drosop- hila subatrata</i>	White cryst. 170-71.	290; end absorption below 230. +20 (c, 3 absol. EtOH, 24°).	Gram +, <i>Myc. smegmatis</i> ; less active against gram—. Influenza PR-8 virus in chick embryo
143*	$C_{22}H_{38}N_2O_6$ Enniatin B	<i>Fusarium orthoceras</i> var enniatum	173-76	—106 to —108 (c, 0.63 to 1.2 CHCl ₃),	Mycobacteria, some fungi
144*	$C_{22}H_{38}O_6$ Ungulmic acid	<i>Polyporus benzoinus</i> , <i>P. betulinus</i>	Microcrystalline needles. Tribasic acid. 78-80.	Opt. inactive.	<i>M. pyogenes</i> var. <i>aureus</i> .
145*	$C_{23}H_{38}O_8$ Ramalinolic acid	<i>Ramalina</i> spp.	Prisms or plates. Depside of orcinol group. 163-64		<i>Staph. aureus</i>
146*	$C_{24}H_{40}O_7$ Anziatic acid	<i>Anzia</i> spp., <i>Cetraria san- guinea</i>	Minute needles. Depside of orcinol group. 124 dec.		<i>Staph. aureus</i>
147	$C_{28}H_{52}O_4$ Ophiobolin	<i>Ophiobolus miyabeanus</i>	White prisms. 181-82.		Fungi
148*	$C_{24}H_{42}N_2O_6$ Enniatin A	<i>Fusarium orthoceras</i> var enniatum	Colourless long needles. 121-22.	—90 to —92 (c, about 1 CHCl ₃).	Gram +, acid fast, fungi
149	$C_{24}H_{42}N_2O_7$ Sambucinin	<i>Fusarium sambucinum</i>	Colourless plates or tetrahe- dra. Neutral 86-87.	Absorption below 219. —83.2 \pm 2 (c, 1 EtOH, 21°)	Gram +, <i>Myc. phlei</i>
150	$C_{24}H_{48}N_2O_6$ Baccatine A	<i>Gibberella baccata</i>	Colourless cryst. 135.		<i>M. pyogenes</i> var <i>aureus</i> , fungi.
151	$C_{25}H_{38}N_2O_6S_2$ Micrococcin	<i>Micrococcus</i> sp.	White, fine, long, silky need- les in fans, sheaves or balls. 222-28 sinters.	345 with blue or purple fluo- rescence. +116 \pm 1 (EtOH, 21°)	Gram +, <i>Vibrio comma</i> , <i>My- co. tuberculosis</i>
152*	$C_{25}H_{28}O_8$ Lobaric acid	<i>Stereocaulon</i> spp., other lichen spp.	Needles. Depsidone or orcinol group 192.		<i>Staph. aureus</i>

1	2	3	4	5	6
153*	$C_{50}H_{122}O_7$ Perlatolic acid	Cladonia spp., Parmelia spp.	Needles. Depside of orcinol group. 108.		Staph. aureus, Myco. tuberculosis avium
154*	$C_{50}H_{122}O_8$ Boninic acid	Ramalina boninensis	Plates. Depside of orcinol group. 134.5.		Staph. aureus
155	$C_{50}H_{144}N_2O_7$ Avenacin	Fusarium avenaceum	Colourless plates or tetrahedra. Neutral. 139.	Absorption below 219. —101 \pm 2 (c, 1 EtOH, 19°).	Gram +, Myco. phlei
156*	$C_{50}H_{130}O_9$ Physodic acid	Parmelia physodes, Parmelia furfuracea	Prisms. Depsidone of orcinol group. 205.		Myco. tuberculosis
157*	$C_{50}H_{132}O_8$ Olivetonic acid	Parmelia olivetorum, Cornicularia spp.	Needles. Depside of orcinol group. 151.		Staph. aureus, Myco. tuberculosis avium
158*	$C_{50}H_{134}O_7$ Fumagillin	A. fumigatus	Colourless or light yellow cryst. Weak monobasic acid. monoester of decatetraenedioic acid. 189-94.	239, 304, 322, 336, 351. —26.6 (c, 0.25 MeOH, 25°); —24 (c, 5 CHCl ₃ , 25°).	Protozoa and viruses
159	$C_{50}H_{144-46}N_2O_7$ Fructigenin	Fusarium fructigenum	Colourless plates or tetrahedra. Neutral. 129.	Absorption below 219. —103 \pm 2 (c, 1 EtOH, 18°).	Gram +, Myco. phlei
160	$C_{50}H_{146}N_2O_7$ Lateritium II	Fusarium lateritium	Colourless plates or tetrahedra. Neutral. 125.	Absorption below 219. —92 \pm 1.6 (c, 1.2 EtOH, 19°).	Gram + Myco. phlei
161	$C_{50}H_{140}O_7$ Bongkrekic acid	Pseudomonas cocovenans	Anorphous.	239, 263 (0.01% NaHCO ₃); 238, 267 (EtOH). +165 (2% NaHCO ₃ , 22°); +105 (96% EtOH, 22°).	Gram +, gram—, fungi.
162	$C_{50}H_{220}O_{10}$ Rugulosin	P. rugulosum, P. wortmanni	Yellow cubes or prisms (from EtOH or acetone). Related to polyhydroxy-anthraquinones. 293 dec.	(∞) $\frac{18}{640}$ + 605 (dioxane; falls to +222 on exposure to light for periods of 14 days or more).	Gram +, gram—, Myco. smegmatis, Pythium intermedium

1	2	3	4	5	6
163*	$C_{30}H_{52}O_2$ Zeorin	Anaptychia spp., lichen spp.	Colourless hexagonal plates. Triterpenoid acid. 253.		Staph. aureus
164*	$C_{31}H_{46}O_4$ Polyporenic acid C	Polyporus benzoinus, P. betulinus	Microcrystals. Triterpenoid acid. 285-90 dec.	$(\alpha)_{546}^{20} + 8.19$ (c, 5.1 pyridine).	Acid fast. Slight activity against M. pyogenes var. aureus and E. coli
165*	$C_{32}H_{42}O_8$ Helvolic acid	A. fumigatus	White, long, fine needles. Monobasic acid. 211.3-212.1; 220-26	230, 234, 231, 322. —113 (c, 3.1 $CHCl_3$, 23°); —117 (c, 2.6 $CHCl_3$, 23°); —124 (c, 1 $CHCl_3$, 25°)	Gram + gram—, acid fast, fungi.
166	$C_{32}H_{48}O_8$ Cephalosporin P ₁	Cephalosporium sp.	Needles. Acidic. Polycyclic aromatic structure. 147.	211—218. +28.	Gram +, Myco. phlei
167*	$C_{32}H_{60}O_{14}$ Glyco-lipidic antibiotic	Pseudomonas aeruginosa	Colourless, thin, rectangular platelets. 86.	—84 (c, 3 $CHCl_3$)	Myco. tuberculosis var. ho- minis.
168*	$C_{35}H_{23-25}N_5O_8$ Violacein	Chromobacterium viola- ceum	Violet black microcryst. Thick needles. Elongated, rectangular cryst. Violet pigment. Above 350 dec.		Gram +, Neisseria meningi- tis, fungi, protozoa
169	$C_{37}H_{62-66}O_7$ Ustilagic acid	Ustilago zeae	Colourless, needle-like cryst. Mixture of closely-related partially acylated glucoli- pids. 141-47.	+7 (c, 1 pyridine, 23°).	Gram, + gram—, fungi, Actinomycetes
170	$(C_{39}H_{74}N_{12}O_9)_n$ Circulin	Bacillus circulans	White amorphous solid (sul- phate). Basic polypeptide. Sulphate: 226-28; HCl: 232-36 dec.	None significant. Sulfate: —61.6 (c, 1.25 water, 25°); HCl: —60.1 (c, 2.094 water, 25°)	Gram—, Less active against gram +, Trichomonas vagi- nalis, fungi.
171	$C_{42}H_{28-30}N_6O_7$ Violaccin (See 168)				
	$C_{48}H_{60}O_{16}$ Glutimosin	Myrothecium verrucaria	Colourless thin plates. Darkens, but no melting by 300.	About + 54 (c, 0.2 C_6H_6 , 20°).	Fungi

1	2	3	4	5	6
172	$C_{50}H_{97}N_{15}O_{15}$, C ₁₄ Polymyxin D (4 HCl)	Bacillus polymyxa	No definite structure; birefringent. Basic polypeptide. 228-35 dec. (for Polymyxins)	No characteristic spectrum.	Gram—
173	$C_{56}H_{106}N_{12}O_{13}$ Polypeptin	Bacillus krzemieniewski	Cryst. form varies with solvent; triangular prisms from 65% EtOH. Basic polypeptide. 176.	252, 258, 264. —93.3 (isopropyl alcohol, 20°).	Gram+, gram—, Myco. tuberculosis, fungi.
174	$C_{59}H_{104}N_{16}O_{14}$, C ₁₅ Polymixin B ₁ (5 HCl)	Bacillus polymyxa	No definite structure; birefringent. Basic polypeptide. 228-35 dec. (for Polymyxins).	No characteristic spectrum. —85.1 (c, 2.33 in 75% EtOH, 25°)	Gram—
175*	$C_{60}H_{92}N_{12}O_{10}$ Gramicidin S	Bacillus brevis	Thin, colourless needles. Cyclic decapeptide. 245-47, 256-58, 268-70.	—292 (c, 1.5 EtOH, 18°); —295 (c, 1.5 EtOH, 20°).	Gram+, gram—
176*	$C_{66}H_{86}N_{13}O_{13}$ Tyrocidine A	Bacillus brevis	Colourless, fine needles or rods (HCl). Basic polypeptide. 240-42.	290. —111 (c, 1.37 in 50% EtOH, 25°).	Gram+, Actinomycetes; less active against gram—, protozoa.
177*	$C_{66}H_{97}N_{15}O_{17}S$ Bacitracin F	Bacillus subtilis; B. licheniformis. Conversion of Bacitracin A	White, amorphous powder. Weakly basic polypeptide.	290.	Gram+
178*	$C_{66}H_{103}N_{17}O_{16}S$ Bacitracin A	Bacillus subtilis; B. licheniformis	White, hygroscopic, amorphous powder. Weakly basic polypeptide.	253. —3.6 (c, 5.0 in 0.01 HAc, 25°).	Gram+
179*	$C_{68}H_{88}N_{14}O_{13}$ Tyrocidine B	Bacillus brevis	Colourless needles or rods (HCl). Basic polypeptide.	As for Tyrocidine A (176).	As for Tyrocidine A (176).
180	$C_{71}H_{112}N_{14}N_{18}O_{17-18}S$ Bacitracin B	Bacillus subtilis	White, highly hygroscopic amorphous powder Weakly basic polypeptide	253. —2.8 (c, 5.0 in 0.01 HAc, 25°).	Gram+
181	$C_{127}H_{169}N_{26}O_{28}$ Tyrocidine (mixture)	Bacillus brevis	HCl: Colourless fine needles or rods. Mixture of basic polypeptides. 237-39 uncor., 240 cor., dec.	—101, —102 (c, 1 in 95% alcohol, 25°).	Gram+, Actinomyces spp., less active against gram—

1	2	3	4	5	6
182	$C_{148}H_{210}N_{30}O_{26}$ Gramicidin	<i>Bacillus brevis</i>	Colourless platelets with pointed or rectangular ends. Natural polypeptide, a component of Tyrothricin. 228-31.	281.5, 290.5, 271. +3 (c, 0.17 in 95% EtOH, 20°)	Gram +, protozoa
ANTIBIOTICS FOR WHICH EMPIRICAL FORMULA IS NOT KNOWN					
183	Xerosin	<i>Achromobacter xerosis</i>	Acid precipitable material.	255-260 (aq. soln.)	In vivo activity in mice infected with influenza A and B viruses, Newcastle disease, mouse pneumonitis
184	Alternarine	<i>Alternaria solani</i>	White needles. 230.		Gram +, gram —, acid fast, fungi.
185	Gigantlin	<i>Aspergillus terreus</i> var. antibioticus	110.		<i>E. coli</i>
186	A. U. N. I	<i>Aspergillus nidulans</i>	Neutral. 155-56.		Gram + cocci and Myco-bacteria.
187	A. U. N. II	<i>Aspergillus nidulans</i>	Neutral. c. 270 dec.		Similar to A. U. N. I (186)
188	A. U. N. III	<i>Aspergillus nidulans</i>	Neutral. 225.5-226.5.		
189	Terrecin	<i>Aspergillus terreus</i>	Light yellow prisms. 219-20.		Gram +, less active against gram—
190	Brevolin	<i>Bacillus brevis</i>	HCl: Yellowish-white, amorphous.	272.5 (99% MeOH). —18.9 (c, 1 MeOH, 26°)	Gram +, some gram—, acid fast
191	Gramicidin A	<i>Bacillus brevis</i>	Neutral polypeptide. 227-228.		Gram +, protozoa
192	Gramicidin B	<i>Bacillus brevis</i>	Neutral polypeptide. 258-59.		Gram +, protozoa
193	Gramicidin C	<i>Bacillus brevis</i>	281.		Gram +, gram—
194	Lichenformins A, B, C	<i>Bacillus licheniformis</i>	HCl: White, amorphous powder. Char on heating. Peptides.	Below 250 (impurities affect u. v. absorption). HCl: A: -37.4; B: -37.7; C: -36.8.	Gram +, gram—. Myco-plei inhibited at higher concn.

1	2	3	4	5	6
195	Esperin	<i>Bacillus mesentericus</i>	Na salt : Needles. 195; Na salt: 268-69 dec.		Myco. tuberculosis
196	Pumilin	<i>Bacillus subtilis</i>	Lemon yellow, small hexagonal crystals. Largely non-peptide. Does not melt below 360.	365, 378, 400.	Gram +
197	Bacillomycin	<i>Bacillus subtilis</i>	Colourless, microscopic needles or rosettes. Polypeptide with titratable acid function.	277 (n-butanol-water-acetic acid-95% ethanol) 296 (alcoholic NaOH) Opt. inactive(EtOH),	Gram +, Gram — at higher concn.
198	Bacitracin C	<i>Bacillus subtilis</i>	White, hygroscopic, amorphous powder. Weakly basic polypeptide.	250, 268.	
199	Bacitracins D, E	<i>Bacillus subtilis</i>	White, hygroscopic, amorphous powder. Weakly basic polypeptide.	253.	
200	Bacitracins F ₁ , F ₂ , F ₃	<i>Bacillus subtilis</i>		253, 288.	
201	Bacitracin G	<i>Bacillus subtilis</i>		250, 268.	
202	Iturin L	<i>Bacillus subtilis</i>	Grey powder. Protein.	278.	Gram +, P. notatum
203	Iturin N	<i>Bacillus subtilis</i>	Yellow powder.	270.	
204	Iturin O	<i>Bacillus subtilis</i>	Brown substance.	260.	
205	Subtenolin	<i>Bacillus subtilis</i>	Light yellow powder, somewhat hygroscopic.	End absorption, 270.	Gram +, some gram —
206	Rhizobacidin	<i>Bacillus subtilis</i>	Crystalline. Peptide. 215-20.		Rhizobium
207	Cephalosporin P ₂	<i>Cephalosporium</i> sp.	White, greasy, amorphous solid. Acidic. 151.	211.	
208	Cephalosporin P ₄	<i>Cephalosporium</i> sp.	Light fawn coloured cryst. Acidic. 220-30.	211.	

1	2	3	4	5	6
209	Cephalothecin	Cephalothecium sp.	124-26 dec.		Fungi.
210	Clitocybins	Clitocybe candida	Colourless, long orthorhombic cryst. 77.		Gram +, gram —, Myco. tuberculosis
211	Diatretyne 3	Clitocybe diatreta	White crystals turning black Neutral.	253, 280, 297, 316, 339. 274 (ether).	Gram +, gram —, Myco. tuberculosis
212	Quadrifidin B ₂	Coprinus quadrifidus	Strongly dichroic white cryst. Neutral.	265, 276, 294, 313 (ether).	Gram +, gram —, Myco. phlei.
213	Quadrifidin B ₃	Coprinus quadrifidus		217, 259, 274, 290, 309 (EtOH).	M. pyogenes var. aureus, B. subtilis.
214	Drosophilin D	Drosophila subatrata	Polyacetylene.	238, 251, 265, 280 (ether).	Gram +, gram —, Myco. smegmatis
215	Drosophilin C	Drosophila subatrata	Polypeptide.	272 broad band.	Fungi and yeast
216	Coliformin	Escherichia coli	I Amorphous black pigment II Yellow pigment. I does not melt or sublime below 360.		W. perfringens
217	Fuligolic acids	Fuligo septica	c. 70.		M. tuberculosis var. hominis
218	Oxysporin	Fusarium oxysporum	Colourless thin needles. 166-68.		Gram +, gram —
219	Lenzitin	Lenzites sepiaria	Colourless orthorhombic plates. 158-60 dec.	No specific absorption	Gram +, Myco. tuberculosis H37Rv
220	Ramycin	Mucor ramannianus	Colourless cryst. 156 dec		Fungi. Weak activity against bacteria
221	Factor	Penicillium albidum	Yellow and brownish rectangular rods. Above 250 dec.		Staph. aureus and Shigella shigae
222	Factor	Penicillium herquei	Yellow or buff-coloured powder. Flavoprotein, glucose-oxidase enzyme.	270-280, 375-380, 450-460 —4.8 (c, 0.012 water, 32°)	Gram +, Gram —,
223	Notatin	P. notatum, other Penicillium spp.			

1	2	3	4	5	6
224	Factor	<i>P. spinulosum</i>	White, fine needles. 183-85.		Grant +, gram—
225	Wortmannin	<i>P. wortmanni</i>	Colourless solid. Neutral 240 dec.		Specific against <i>Botrytis alli</i> , <i>B. chereca</i> , <i>B. fabae</i> , <i>Glad-</i> <i>osporium herbarum</i> , <i>Rhi-</i> <i>zopus stolonifer</i> .
226	Biformyne 1	<i>Polyporus biformis</i>	Polyacetylene.	259, 274, 291, 310.	Gram +, gram—, acid fast, fungi.
227	Biformyne 2	<i>Polyporus biformis</i>	Polyacetylene.	263, 274, 290, 310.	Similar to that of <i>Biformyne</i> 1 (226)
228	Comirin	<i>Pseudomonas antimycetica</i> , <i>Bacterium antimyceticum</i>	Whitish amorphous powder Peptide containing an ether soluble moiety. 230-35 dec. depending on rate of heating.		Fungi and yeast
229*	Viscosin	<i>Pseudomonas viscosa</i>	Acidic polypeptide. 269 dec., 270-73 dec.	—162.2 (20°), 168.3 (29°)	Myco. tuberculosis
230	Marcescin	<i>Serratia marcescens</i>	Polypeptide.	240, 260.	Gram +, gram—, Myco. tuberculosis
231	Sphaerophorin	<i>Sphaerophorus compressus</i>	140.		<i>M. pyogenes</i> var. <i>aureus</i>
232	Nisins A, B, C, D	<i>Streptococcus crenoris</i> , <i>Streptococcus lactis</i>	Nearly white needles. Low molecular weight pro- tein or polypeptide.	275.	Gram +, gram—, Myco. tuberculosis, Actinomycetes.

MANUFACTURED FROM THE BASIC STAGE

FOR THE FIRST TIME IN INDIA

PROCAINE

HYDROCHLORIDE B.P.

HOECHST


**PHARMACEUTICALS
PRIVATE LIMITED**

Backbay Reclamation, Bombay-I



Manufactured in collaboration with
FARBWERKE HOECHST A.G.
Frankfurt (M) – Hoechst/Germany

HF. 3-432



*The new I.C.I. Pharmaceutical
Research Laboratories, Cheshire*

I.C.I. Leadership in Pharmaceutical Research

In I.C.I.'s new laboratories at Alderley Park, Cheshire, chemists, biologists, doctors, pharmacists and veterinary surgeons are working on a continuous, well-coordinated programme for the conquest of disease with new and better drugs. Already the I.C.I. laboratories have made a number of notable discoveries which have played a major role in protecting the health of communities all over the world.

Some recent notable I.C.I. contributions in human medicine are :

'PALUDRINE' • 'MYSOLINE' • 'HIBITANE'
'CETAVLON' • 'FLUOTHANE'

These, together with other important therapeutic agents produced by I.C.I., are to be found in the comprehensive range of medical and veterinary products marketed by Imperial Chemical Industries (India) Private Limited.

IMPERIAL CHEMICAL INDUSTRIES (INDIA)
PRIVATE LIMITED

Calcutta Bombay Madras New Delhi



ICP 8/1



With Compliments from

MINIMAX LIMITED

(FIRE PROTECTION ENGINEERS)

SUPPLIERS OF

INDIGENOUSLY MANUFACTURED FIRE
EXTINGUISHERS - & TRAILER - PUMPS.

FREE AFTER SALES SERVICE A SPECIALITY

PLEASE ENQUIRE :

MINIMAX LIMITED

29, FORT STREET,

BALLARD ESTATE,

BOMBAY 1.

THE SIGN OF
SECURITY !

Tele. Office : 262879

Tele. Workshop : 76167

DISTRIBUTORS OF DISTINCTION :

- **LEDERLE LABORATORIES (INDIA) PRIVATE LTD., BOMBAY**
(For Northern India)
- **HINDUSTAN ANTIBIOTICS LTD., PIMPRI**
- **CAPCO PRIVATE LTD., E. MERCK DEPARTMENT, GERMANY**
(For Northern India)
- **KESAVARDHINI PRODUCTS, COIMBATORE**
(For Northern & Western India)
- **HYGIENIC RESEARCH INSTITUTE, BOMBAY**
VASMOL HAIR DARKENER & TONIC
FLOROZONE BEAUTY CREAM & LOTION
- **DIAL RUBBER WORKS, BOMBAY**
SURGICAL GLOVES
- **FAMILY PLANNING (WHITTAKE LABORATORIES, INC., U.S.A.)**
- **DENTAL EQUIPMENT AND SUNDRIES, KOVO & CHEMAPOL,**
CZECHOSLOVAKIA

Dr. Jai Singh's Son and Co. Private Ltd.

18/4, ASAF ALI ROAD, NEW DELHI.

NEW INDIA DEVELOPMENT CORPORATION (PRIVATE) LTD.

OFFER TO SUPPLY FROM THEIR
MODERN WORKS AT PIMPRI (POONA)

SUGAR MACHINERY

MANUFACTURED UNDER SUPERVISION OF GERMAN
EXPERTS AND IN TECHNICAL CO-OPERATION WITH



BUCKAU-WOLF
OF WEST GERMANY



ORIGINAL DESIGNS, DRAWINGS AND PATENTS
ARE BEING USED

Please send your enquiries to the
SOLE SELLING AGENTS

PROTOS ENGINEERING CO. (PRIVATE) LTD.

173, Jamshedji Tata Road, Churchgate, Bombay.

Telegrams: "PROTODYN"

Telephone: 37147-8-9

BOMBAY
P. O. Box 1642

CALCUTTA
P. O. Box 2549

DELHI
P. O. Box 431

MADRAS
P. O. Box 375



Naturally, now that DET, Swastik's modern detergent, makes your clothes a dream of cleanliness. You know DET is equally good in hard or soft, warm or cold water. You can even clean floors, sinks, crockery, cutlery, etc. gleaming bright with DET.

Swastik Oil Mills Ltd., Bombay



Shilpi S.O.M.66



The Priceless Ingredient

In the city of Bagdad lived Hakeem, the Wise One, and many people went to him for counsel, which he gave freely to all, asking nothing in return.

There came to him a young man, who had spent much but got little, and said: "Tell me, Wise One, what shall I do to receive the most for that which I spend?"

Hakeem answered: "A thing that is bought or sold has no value unless it contains that which cannot be bought or sold. Look for the Priceless Ingredient."


"But, what is this Priceless Ingredient?" asked the young man.

Spoke then the Wise One: "My son, the Priceless Ingredient of every product in the market-place is the Honor and Integrity of him who makes it. Consider his name before you buy."

Copyright 1955 E. R. Squibb & Sons division of Olin Mathieson Chemical Corporation

SQUIBB

SARABHAI CHEMICALS BARODA
Karamchand Prichand Private Limited
MANUFACTURERS OF SQUIBB MEDICINALS IN INDIA



GLAXO LABORATORIES (INDIA) PRIVATE LTD.

Manufacturers of quality Pharmaceuticals,

Fine Chemicals, Antibiotics,

Immunologicals and Infant Foods.



Parry's

DISTRIBUTORS FOR



Antibiotics: Haematinics:
Chemotherapeutics: Vitamins:
(Distributors in Southern & Eastern India)



Penicillin &
Streptomycin products.

Govt. Oil Factory,
Kozhikode.

Sea Gold Shark Liver Oil.
Stayfit Shark Liver Oil & Capsules.
Adamin Liquid & Capsules.

Madras Govt.
Quinine Factory.

Quinine Products Eucalyptus Oil
(Distributors for Southern & Western India)

Johnson & Johnson

Toiletries, Surgical Adhesive Tapes.
Bandages and Dressings and ORTHO Products.
(All India except Bombay, Madhya Pradesh
& Rajasthan States)



Surgical Ligatures. Catguts:
Mersutures:
Mersilk: Mersilene.



Absorbent Cotton Wool,
Bandage: Gauze: Lint.



Operating & Diagnostic
Surgical Instruments.
(Distributors for
Southern India)

PARRY & CO. LTD.,

MADRAS, Post Box No. 12, • CALCUTTA Post Box No. 208,
BOMBAY, Post Box No. 506, • NEW DELHI Post Box No. 173.

G. PPII,32